

Effects of the COVID-19 Pandemic on Physical Activity in Children: A Systematic Review

Alex Ramirez,¹ Ashley B. Rapp,² Sara Santarossa.³

Abstract

Background: Extended lockdowns established to minimize the spread of COVID-19, may have affected physical activity (PA). The objective of this systematic review is to report how PA of children was impacted by the COVID-19 pandemic. **Methods:** A database search of CINAHL, Embase, MEDLINE, PsycInfo, and Web of Science was conducted on 03/29/2021 in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines. Studies were included if they reported a measure of PA for children ≤ 18 years and were published in English in a peer-reviewed journal. The search generated 673 unique citations. After applying the criteria above, 69 articles were included. The quality of the included articles was assessed using the NIH NHLBI Study Quality Assessment Toolbox. **Results:** Unstructured play, walking, and virtual PA (e.g., YouTube, online classes) were the most frequently reported modes of PA during the COVID-19 pandemic. The PA of children decreased among 89% of articles that reported changes in PA (e.g., frequency, intensity, duration, or the percentage of children who met PA recommendations) of children during the COVID-19 pandemic. Increased PA was reported more often among younger children, boys, children who lived with other children, and children with more outdoor space. **Conclusion:** Globally, PA of children decreased during the COVID-19 pandemic. Further work is needed to develop policies to support global PA increases. Limitations of this review include the use of online surveys which are limited by participants ability to recall PA behaviors and access to online resources.

Introduction

The COVID-19 pandemic (the pandemic) and its effects rapidly permeated the globe with over 100 million cases by February 1st, 2020.¹ By March 2020, many governments issued stay-at home orders, shelter-in-place orders, lockdowns, and various social-distancing guidelines to mitigate the spread of COVID-19.² These restrictions, including the closure of schools, parks, and organized sports, may have altered movement behaviors, including physical activity (PA).^{3,4} PA in children encompasses a variety of modes (e.g., exercise, sports, and active play).⁵ It has been defined as any body movement that increases energy expenditure and can be characterized by the components of the FITT principle: frequency, intensity, time (e.g., duration), and type.⁶ The WHO recommends children (ages 5-17) engage in ≥ 60 min/day of moderate-to-vigorous (MVPA) intensity PA⁷ where 3 days/week include vigorous-intensity aerobic activities, muscle-strengthening and bone-strengthening activities.⁸ As a critical component of child development, PA has substantial long-term health benefits including the prevention of chronic health conditions.^{9,10} Conversely, insufficient PA in children can increase the risk of developing cancer, heart disease, obesity, and type 2 diabetes.¹¹ Despite the known benefits of PA on health, global levels of PA were insufficient in 4 out of 5 children prior to the pandemic.¹² Thus, the prolonged stay-at-home orders brought on by the pandemic may have exacerbated the proportion of children with

insufficient PA, as previous studies have shown that children have less PA when they are outside of school (e.g., holidays, summer, weekends).^{13,14} Although children seem to have a lower risk of contracting COVID-19,^{15,16} the physical health burden of the pandemic and the resulting restrictions are of increasing interest.^{1,17,18}

Previous studies have demonstrated that childhood health behaviors, including PA, are likely to persist through adulthood and can impact lifetime health.^{19,20} It is important to monitor PA changes in children to reduce the risk of developing long-term health consequences associated with inadequate PA (e.g., cardiovascular disease, type 2 diabetes, obesity).^{21,22} Not only can PA prevent many chronic diseases,²³ but it can also have acute health benefits such as boosting the immune system.^{24,25} A recent study in adults infected with COVID-19 found reduced risk of critical outcomes in individuals who routinely adhered to PA recommendations.²⁶ Although this relationship has not yet been established in children, several studies have found obesity to be the most prevalent characteristic in children infected with COVID-19.^{27,28} The disruption of PA in children may increase the prevalence of childhood obesity and other comorbidities which could, in turn, increase the risk of infection and subsequent severity in outcomes from COVID-19.^{15,29}

¹ B.S., Third-year Medical Student, Wayne State University School of Medicine/Henry Ford Health System, Detroit, USA.

² MPH. Henry Ford Health System, Detroit, USA.

³ PhD. Henry Ford Health System, Detroit, USA.

About the Author: Alex Ramirez is currently a third-year medical student at Wayne State University School of Medicine in Detroit, MI, USA of a 4 year program. She has been conducting research in the Public Health Sciences Department at Henry Ford Health since 2020.

Correspondence:

Alex Ramirez

Address: 540 E. Canfield Ave. Detroit, MI 48201, USA.

Email: alex.ramirez@med.wayne.edu

Editor: Francisco J. Bonilla-Escobar

Student Editors: Shuo-Yan Gau, Diego Carrion Alvarez

Copyeditor: L V Simhachalam Kutikuppala

Proofreader: Laeeqa Manji

Layout Editor: Julian A. Zapata-Rios

Submission: Sep 22, 2022

Revisions: Feb 8, 2023

Responses: Feb 9, 2023

Acceptance: Jun 11, 2023

Publication: Jun 13, 2023

Process: Peer-reviewed

As childhood is a critical period of development, understanding the implications of the pandemic on health behaviors is important and there is a need to highlight potential adverse effects that may need to be actively monitored and remediated as COVID-19 restrictions lift and vaccines are distributed. Identifying changes in PA (FITT principle) of children is important for understanding the extent to which the pandemic impacted the already low levels of global PA among children. In addition, recognizing the biological, social, and environmental factors that affected changes in PA is critical for the development of effective programs to prevent short- and long-term health consequences. To the authors' knowledge, this is the first review to synthesize the literature at the time of the data search regarding the global impact of the pandemic on the duration and types of PA children participated in and what biological, social, and environmental factors contributed to those changes. The goal of this review was to synthesize the global impact of the pandemic on PA of children by addressing the following questions:

- 1) What types of PA did children participate?
- 2) To what extent did PA (e.g., duration of PA, percentage of children meeting PA recommendations) of children change?
- 3) What biological, social, and environmental factors affected changes in PA of children?

Methods

This review was conducted using the PRISMA guidelines.³⁰ The selection process for the included articles is summarized in [Figure 1](#).

Search Strategy

Two research librarians independently searched MEDLINE, Embase, PsycInfo, CINAHL, and Web of Science on 03/29/2021. The searches were limited to the English language, COVID-19, and 2020-current (03/29/2021). The key words used in the search strategy included various combinations of terms such as: "physical activity," "exercise," "activity level," or "sports" AND "child," "adolescent," "girl," "boy," or "youth" AND "covid-19," "severe acute respiratory syndrome coronavirus 2" or "sars-cov-2". Following their independent searches, the librarians compared results and removed duplicates.

Article Inclusion and Exclusion Criteria

Included studies were those that were published in English in a peer-reviewed journal, measured PA during the pandemic, and encompassed PA data for children ≤ 18 years. Articles were excluded if they did not present original empirical findings, if they described the development of a tool or measure, or if they were inaccessible to the authors (e.g., no free full-text download was available).

Article Selection


After the librarians eliminated duplicate articles, 673 articles remained. The following article selection process was conducted in 4 steps. First, 2 authors conducted an independent abstract review to determine whether each article should be included or

not based on the inclusion and exclusion criteria. Second, the 2 authors discussed any articles they disagreed on until consensus. Following this discussion, 572 articles were excluded because they had a study population outside the context of the current review, meaning that PA data presented in the findings were not specified for participants ≤ 18 years old. Third, the full text of the remaining 101 articles was reviewed to ensure the articles were within the scope. The data extracted included authors, year of publication, sample source (e.g., the country the sample was gathered from), sample size, sample age range, PA measure, main findings, and factors (e.g., biological, social, and environment) related to PA and are outlined in [Table 1](#). Ultimately, 32 of these articles were excluded because they met ≥ 1 exclusion criteria. In the fourth step, all authors met to confirm the final 69 articles should be included. The data from the 69 articles was synthesized into a table adapted from a review paper of a similar topic [Table 1](#).³¹

Quality Assessment

The quality of the included articles was assessed using the NIH NHLBI Study Quality Assessment Toolbox.³² This quality assessment tool describes a "good" study as one with a low risk of bias and a "fair" study as one with some bias.³²

Figure 1. Article Selection Diagram Using PRISMA.



Results

General Characteristics

General characteristics of the included articles are shown in [Table 2](#). Most articles included participants in Europe (n=34),³³⁻⁶⁶ North America (n=13),⁶⁷⁻⁷⁹ or South America (n=6).^{49,58,80-83} The distribution of countries represented by the articles in this review are depicted in a heat map in [Figure 2](#).

Table 1. Characteristics of Included Articles (N=69) and Effects of the COVID-19 Pandemic on Physical Activity (PA) and Biological, Environmental, and Social Factors that Affected PA.

No.	Author (Year)	PA Measure; PA Data Reported by ^a	Sample Source (Country)	Sample Size	Age Ranges	Findings	Biological, Environmental, and Social Factors that Affected PA	NIH NHLBI Quality Assessment
33	Alonso-Martinez (2021)	Wrist-worn GENEActiv tri-axial accelerometer	Spain	Pre-pandemic: 268 children. Pandemic (03-04/2020): 145 children, PA measured in 21	4-6	On average, preschoolers had 43.3 minutes less/day (95% confidence interval (CI) -68.1 to -18.5 (p=.002)) of total PA during the Pandemic. MVPA also decreased an average of 17 min/day (p<0.001).	Children who met PA recommendations had lower internalizing scores (i.e. social withdrawal, shyness, anxiety).	Good
34	Androutsos (2021)	Online survey; 2	Greece	397 children	2-18	During lockdown, 66.9% of parents reported their child's PA level decreased.	Decreased PA was associated with body weight increase (p<0.001).	Good
35	Cachon-Zagalaz (2021)	Online survey; 2	Spain	837 children	0-12	The highest percentage of participants (34%) spent 0 days/week on PA during lockdown, followed by 32.6% for 2-3 days/week, 19.6% for 4-5days/week, and 13% for 6-7 days/week.	Children with an established routine/schedule (M = 38.13; SD = 35.01) and children in the 6 to 12 years old age group (M = 38.95; SD = 30.25) reported higher levels of daily PA. 34.8% of the sample reported 0 days/week of PA during the Pandemic.	Good
36	Chambo-nniere (2021)	Online survey; 1,2	France	6491 children	6-17	During lockdown, PA decreased for 42% of children, stayed the same for 21.3% of children, and increased for 36.7% of children. Similarly, PA decreased for 58.7%, stayed the same for 21.8%, and increased for 19.6% of adolescents.	PA decreased among 35.2% of children in rural areas, 46.7% of children in suburban areas, and 47.9% of children in urban areas (PA was significantly associated with geographic location p<0.001). Living in an urban environment was associated with a decrease in PA (p<0.001). Among children without access to an outdoor area, 64.2% reported decreased PA, while among children with access to an outdoor area, only 37.8% reported decrease PA during the Pandemic (p<0.001). The proportion of participants who increased, decreased or did not change their PAL during the lockdown was significantly associated with gender in adolescents (p<0.001) but not in children (p=0.10). Children and adolescents who were more active prior to the Pandemic had greater reductions in PA compared to those who were initially less active (p>0.001).	Good
37	Delisle (2020)	Telephone questionnaire; 2	Sweden	100 children	3-5	PA (+53 min/day) and outdoor time on weekdays (+124 min/day) and weekends (+68 min/day) significantly increased (all p-values ≤ 0.001). PA guidelines were met by 90.3% (n = 65) of the children.	Boys had higher levels of PA (262.5 ± 60.3 vs. 220.4 ± 39.2 min, p = 0.002), MVPA (136.9 ± 43.8 vs. 107.8 ± 29.2 min, p = 0.004), and vigorous PA (39.0 ± 19.4 vs. 29.4 ± 15.7 min, p = 0.028) compared to girls. However, gender was not significantly associated with meeting PA recommendations.	Good
38	Gilic, B., et al. (2020).	PAQ-A; 1	Bosnia and Herzegovina	688 children	15-18	Compared to before the Pandemic, the PAL of the entire sample decreased (from 2.98 ± 0.71 to 2.31 ± 0.68; t-test: 11.88, p < 0.001), as well as among girls (2.69 ± 0.49 to 1.95 ± 0.56; t-test: 2.50±0.44; t-test: 10.01, p<0.01).	Boys were more likely to achieve a sufficient PAL during the Pandemic (OR: 2.41, 95%CI: 1.11–4.01). Higher paternal education was associated with a greater likelihood of achieving a sufficient PAL during the Pandemic (OR: 1.33, 95%CI: 1.19–2.01). In contrast, children with more parental/family conflict had a lower likelihood of achieving a sufficient PAL (OR: 0.77, 95%CI: 0.60–0.99).	Good

No.	Author (Year)	PA Measure; PA Data Reported by ^a	Sample Source (Country)	Sample Size	Age Ranges	Findings	Biological, Environmental, and Social Factors that Affected PA	NIH NHLBI Quality Assessment
39	Gilic, B., et al. (2021).	PAQ-A; 1	Bosnia and Herzegovina	661 children	15-18	67% of boys and 28% of girls (48% overall) had a sufficient PAL before the Pandemic, while 37% of boys and 9% of girls (24% overall) had a sufficient PAL during the Pandemic (April 2020).	Children with a sufficient PAL during the Pandemic had higher participation in individual and team sports prior to the Pandemic (MWZ: 6.65, 7.95, respectively, all $p < 0.01$).	Good
40	Sekulic, D., et al. (2020).	PAQ-A; 1	Croatia	388 children	15-18	There was a significant decrease in PAL of children from before to during the Pandemic (2.99 ± 0.70 and 2.67 ± 0.60 for PAL-BL and PAL-FU, respectively; t -test = 3.46, $p < 0.001$).	The decrease in PAL of boys (3.10 ± 0.78 and 2.79 ± 0.82 for PAL-BL and PAL-FU, respectively; t -test = 5.15, $p < 0.001$), but not girls (2.71 ± 0.66 and 2.59 ± 0.90 , respectively; t -test = 0.61, $p > 0.05$) was statistically significant. Boys had significantly higher PALs than girls during the Pandemic (t -test = 2.11, $p < 0.05$), with	Good
41	Zenic, N., et al. (2020).	PAQ-A; 1	Croatia	823 children	14-18	PAL decreased among the entire sample (from 2.97 ± 0.61 to 2.63 ± 0.68 , $p < 0.01$) as well as for urban adolescents (from 3.11 ± 0.64 to 2.68 ± 0.67 , $p < 0.001$).	Children who lived in urban areas had greater reductions in PA level.	Good
42	Francisco, R., et al. (2020).	Online survey; 2	Italy, Spain, and Portugal.	1480 children	3-18	During the Pandemic, most children experienced < 30 min of PA daily (53%) compared to 30 to 60 minutes daily before the Pandemic (33.1%). The changes in PA from before to during quarantine were significant for the entire sample ($z = -25.56$, $p < 0.001$, $r = 0.66$) as well as for each country (Italy: $z = -16.08$, $p < 0.001$, $r = 0.60$; Spain: $z = -15.45$, $p < 0.001$, $r = 0.74$; Portugal: $z = -12.48$, $p < 0.001$, $r = 0.66$).	Not reported.	Good
43	Hommes, F., et al. (2021).	Questionnaire; 1	Germany	385 children	8-18	42.2% of primary school students (8 to 13 years old) reported decreased PA during the Pandemic.	A greater proportion of primary school students (8 to 13 years old) reported decreased PA than secondary school students (13 to 18 years old) during the Pandemic.	Good
44	Konstantinou, C., et al. (2021).	PAQ-C; 2	Greece	1509 children	5-14	Compared to before the Pandemic, children's PA decreased in school and out of school after schools were re-opened ($p < 0.001$). The frequency of activities (i.e., sports, cycling, dance) children engaged with also decreased compared to before the Pandemic (post-Pandemic: median [Q1, Q3]: 1.38 [1.25, 1.62] vs. pre-Pandemic: 1.5 [1.38, 1.75]).	Not reported.	Fair
45	Kovacs, V. M., et al. (2021).	PAQ-C; 1, 2	Russian Federation, Spain, Italy, Germany, France, Belgium, Portugal, Romania, Hungary, Poland, and Slovenia	8395 children	6-18	Among the entire sample, children met PA guidelines an average of 4 days/week and 19% of children met the 60 minutes of moderate-to-vigorous PA/day guideline during the Pandemic. However, most countries, individually, had $< 20\%$ of children able to meet PA recommendations (range of 7.5% to 26.7%). Slovenia and Romania had the greatest proportion of children able to meet PA recommendations (26.7% and 23.5%) as well as the greatest proportion of children playing outside for more than 2 hrs/day (56.1% and 66.3%). During the pandemic, 56.6% of children were active in online P.E.	In countries mildly affected by the Pandemic (Germany, Romania, Poland, Slovenia, and Hungary), children were more likely to meet PA recommendations if they had a structured daily routine (OR = 1.62 [95%CI, 1.24–2.13]). Children were also more likely to meet PA recommendations if they played outdoors for more than 2 hrs/day, however this relationship was more predominant among older children (OR = 2.56 [95%CI, 1.98–3.32]). Among older children in countries strongly affected by the Pandemic (Spain, Italy, France, Russia, Portugal), children who participated in online P.E. were more likely to have higher levels of PA (OR = 1.27 [95%CI, 1.12–1.44]).	Good

No.	Author (Year)	PA Measure; PA Data Reported by ^a	Sample Source (Country)	Sample Size	Age Ranges	Findings	Biological, Environmental, and Social Factors that Affected PA	NIH NHLBI Quality Assessment
46	Lopez-Bueno, R., et al. (2021).	VO2 (20m shuttle run)	Spain	89 children	12-14	The average VO2 max of the sample decreased 0.5 ml.kg ⁻¹ .min ⁻¹ (SD 0.3) (p = 0.12) from before to after the Pandemic (before: 46.2 ml.kg ⁻¹ .min ⁻¹ (SD 0.6), after: 45.7 ml.kg ⁻¹ .min ⁻¹ (SD 0.7)). The greatest reduction in VO2 max was reported for 14-year-old girls, with a reduction of 1.5 ml.kg ⁻¹ .min ⁻¹ (SD 0.6) (p=0.02).	Not reported.	Good
47	Lopez-Bueno, R., et al. (2020).	Online survey; 2	Spain	860 children	3-16	PA decreased for all gender and age groups from before to during the Pandemic, with an average of 96.1 minutes of PA/week during the Pandemic compared to 198.6 (SD 180.9) minutes of PA/week before (-102.5 minutes/week). The greatest reduction in PA was among 6- to 12-year-old children with a reduction of 120.4 (SD 159.0) min/week.	Boys and younger children participated in the most weekly minutes of PA (211.9 (SD 188.4) and 223.0 (SD198.0) min/wk).	Good
48	Medrano, M., et al. (2021).	"The Youth Activity Profile" questionnaire (YAP); 1	Spain	113 children	8-16	Children participated in PA an average of 63 (SD 39) min/day during the Pandemic, with an average decrease in PA of 91 ± 55 min/day (P < .001) compared to before the Pandemic. Decreased PA was reported among 95.2% of children during the Pandemic.	Children of mothers with a higher education level had smaller reductions in PA during the Pandemic (-1.3 ± 0.1 hour/day vs -1.7 ± 0.1 hour/day, P < .005). Children with access to an outdoor area at home or a large indoor space (i.e. attic or garage) also had lower reductions in PA compared to children without access. (-1.4 ± 0.1 h/d vs -1.7 ± 0.1 h/d, P < .01).	Good
49	Lopez-Gil, J. F., et al. (2021)	Online survey; 2	Spain and Brazil	1099 children	3-17	Compared to before the Pandemic, Spanish and Brazilian children engaged in less days of PA/week during the Pandemic (p<0.001, p<0.001). Before the Pandemic, 34.6% of Spanish children and 30.6% of Brazilian children met PA recommendations. During the Pandemic, 26.5% of Spanish children and 21.8% of Brazilian children met PA recommendations.	Not reported.	Good
50	Morgul, E., et al. (2020).	Online survey; 2	UK	927 caregivers	5-11	Daily PA of children significantly decreased during the Pandemic (x2 (15, n=927) = 121.26, p < .001). Before the Pandemic, 67.3% of children met PA recommendations (at least 60 min of PA/day), while 51.1% of children met PA recommendations during the pandemic.	Not reported.	Good
51	Poulain, T., et al. (2021).	Online questionnaire; 2	Germany	285 children	1-10	From before the during the Pandemic, there was a significant decrease in indoor sports (p=0.001) and an increase in the frequency of outdoor play (before 60%, during 71%).	Lower SES was associated with less frequent outdoor play, while more frequent outdoor play was significantly associated with more children at home (p<0.001).	Fair
52	Ng, K., et al. (2020).	PACE +; 1	Ireland	1214 children	12-18	During the Pandemic, 50% of children reported decreased PA, 30% reported no change in PA, and 20% reported increased PA compared to before the Pandemic. The most frequently reported barriers to PA included coronavirus and sports cancellation, while facilitators to PA included having nothing else to do, school cancellation, and going on walks.	Children with well-established PA habits before the Pandemic were less likely (OR=0.4, CI=0.2-0.6) to report decreased PA, while overweight (OR=1.8, CI=1.2-2.7) or obese (OR=2.2, CI=1.2-4.0) children were more likely to report decreased PA during the Pandemic. Furthermore, participation in strength-training exercises a minimum of three times in the past week was strongly associated with higher levels of PA (OR=1.7, CI=1.3-2.4).	Good

No.	Author (Year)	PA Measure; PA Data Reported by ^a	Sample Source (Country)	Sample Size	Age Ranges	Findings	Biological, Environmental, and Social Factors that Affected PA	NIH NHLBI Quality Assessment
53	Orgiles, M., et al. (2020).	Online survey; 2	Italy and Spain	1143 parents	3-18	During lockdown, the time children spent doing PA decreased ($\beta = 0.04$, SE = 0.07, Wald χ^2 95% CI [0.03, 0.04], $p < 0.001$). For example, the proportion of children who reported < 30 minutes of PA/day increased from 13.6% before the Pandemic to 55.6% during the Pandemic. Similarly, the proportion of children who met PA recommendations (at least 60 mins of PA/day) decreased from 54.1% to 14.8%.	Easier family coexistence was associated with a higher duration of exercise in children ($\rho = 0.08$; $p = 0.004$). In contrast, children of parents with high stress levels were more likely to have less PA ($\rho = -0.10$; $p \leq 0.001$).	Good
54	Pietrobelli, A., et al. (2020).	Telephone questionnaire; 2	Italy	41 children	6-18	Among kids with obesity, sports activity decreased by 2.30 ± 4.60 hours/week ($p=0.003$).	Not reported.	Good
55	Pombo, A., et al. (2020).	Online survey; 2	Portugal	2159 children	0-12	During the Pandemic, 0 to 2 year old children had the highest percentages of PA (%PA). (%PA: play with PA (tag, hide and seek, etc.) + PA (organized indoor and outdoor PA) / (intellectual activity + playful screen time + play w/o PA + play w/ PA + PA)).	Outdoor space ($p < 0.001$), the presence of other children in the home ($p=0.002$), younger age ($p < 0.001$), and having an adult at home not working were significantly associated with higher levels of %PA ($P < .001$). In contrast, lower levels of %PA in children were associated with families in which all adults worked from home ($p=0.016$). There was no significant effect of gender on differences in %PA.	Good
56	Predieri, B., et al. (2020).	Telephone questionnaire; 1	Italy	62 children	1-18	Among children with type 1 diabetes, time spent on exercise significantly decreased from before to during the Pandemic (3.27 ± 2.82 vs. 0.24 ± 0.59 h/week, respectively; $p < 0.0001$).	Not reported.	Good
57	Roe, A., et al. (2020).	Online survey; 2, 3	Norway	5368; 4642 parents, 726 teachers	6-16	About one-third of all participants reported their child participated in more than 60 minutes of daily PA. Students in grades 1-4 (5 to 10 years) had the most PA, with more than 60 minutes of daily PA reported by 43% of parents. In contrast, only 18% of students in grades 8-10 (13 to 16 years) reported more than 60 minutes of daily PA.	Duration of daily PA increased with age of participants. Boys participated in more PA than girls in grades 1-7.	Good
58	Ruiz-Roso, M. B., et al. (2020).	IPAQ; 1	Italy, Spain, Brazil, Chile, and Colombia	726 children	10-19	Before the Pandemic, 27% of all participants were physically active compared to 20.5% during the Pandemic.	Boys were more active than girls [OR 2.22 (CI 95% 1.28–3.86)] before and during the Pandemic. Higher maternal education was associated with lower levels of PA during the pandemic [OR 0.40 (CI 95% 0.20–0.84)].	Good
59	Schmidt, S. C. E., et al. (2020).	MoMo PAQ; 1	Germany	1711 children	4-17	While the duration of organized and nonorganized sports decreased (10.8 min/day, $p < 0.01$), there was an increase in habitual PA (i.e., playing outside, walking, cycling, gardening, housework) (36.2 min/day, $p < 0.01$). The proportion of children who met PA recommendations increased by 11.1% ($p < 0.01$) during the Pandemic. Among the entire sample, the number of active days also increased by 0.4 days/week ($p < 0.01$).	Older children (ages 14 to 17) had greater increases in nonorganized sports than younger children (ages 4 to 5) (17.0 vs 11.8 min/day). Older children had greater reductions in total sports than younger children (15.6 vs 2.2 min/day). Younger children had higher levels of habitual PA overall, but also played outside more, while older children walked and cycled more. Boys reported more min/day playing outside, walking, and cycling than girls. The proportion of children who met PA recommendations increase among 14.7% of children aged 4 to 5 compared to only 4.8% of children aged 14 to 17.	Fair
60	Schmidt, T. and C. S.	Online survey; 1	Denmark	142 children	15-18	From before to during the Pandemic, the average minutes of PA/week decreased by 36.6% ($p < 0.001$). Prior to the Pandemic, most children reported	Not reported.	Good

No.	Author (Year)	PA Measure; PA Data Reported by ^a	Sample Source (Country)	Sample Size	Age Ranges	Findings	Biological, Environmental, and Social Factors that Affected PA	NIH NHLBI Quality Assessment
	Pawlowski (2020).					being physically active 2 to 4 times per week (50.4%), while during the Pandemic most children reported being physically active 5 times or more per week (44.9%). The proportion of children who participated in PA 5 times or more per week increased by 19.1%, while the proportion of children who rarely or never did PA increased by 71.8%. Although outdoor activity was still a frequently reported form of PA, less children reported doing street activities and team ball games compared to before to Pandemic. 57.1% of children reported doing the most PA at home and 33.3% reported doing the most PA in public. Children were most physically active with family (40.8%) or alone (57.7%) in comparison to with friends (83.5%) before the pandemic.		
61	Salzano, G., et al. (2021).	Online survey; 1	Italy	1860 children	12-18	During the Pandemic, 84.5% of children participated in PA at home. PA at home was most often practiced for 1 to 3 hrs/week (30.1%), followed by < 1 hour per week (19.1%).	Not reported.	Fair
62	Ten Velde, G., et al. (2021).	Cohort A: BAECKE questionnaire; Cohort B: BAECKE questionnaire and ActiGraph GT3X accelerometer; 1	Netherlands	Cohort A: 102 children; Cohort B: 131 children	Cohort A: 4-18; Cohort B: 7-12	During the Pandemic, decreased PA was reported by 62% of children in cohort A and 54% of children in cohort B. In cohort A, school, sports, and leisure time PA significantly decreased ($p < 0.01$) compared to before the Pandemic, whereas in cohort B, only sports PA significantly decreased ($p < 0.01$). In cohort B, light PA (LPA) and MVPA decreased from 252 ± 34 and 65 ± 18 min/d before the Pandemic (May/June 2019) to 218 ± 39 and 48 ± 18 min/d during the Pandemic (June 2020) (-34 ± 42 min/d, $p < 0.01$; -17 ± 19 min/d, $p < 0.01$). Similarly, while 64% of children in cohort B met PA recommendations (60 minutes of MVPA/day) before the Pandemic, only 20% met recommendations during the Pandemic ($p < 0.01$). Total PA in cohort B was decreased by 51 ± 52 min/d compared to before the pandemic.	While girls had larger decreases in light PA (LPA) and MVPA than boys, only the difference in reduction of LPA between boys and girls was significant (-18 ± 38 vs -44 ± 41 min/day, $p = 0.02$). Children with more MVPA min/d before the Pandemic had larger reductions in MVPA compared to children with less MVPA before the Pandemic.	Good
63	Theis, N., et al. (2021).	IPAQ-SF; 2	UK	125 parents	8-16	The frequency of vigorous and light/moderate intensity PA decreased from 2 and 5 to 0 and 3 days/week from before to during the Pandemic (median values; $z = -4.699$, $P < 0.001$; median values; $z = -3.979$, $P < 0.001$). However, the duration of vigorous and light/moderate intensity PA did not significantly change with most children achieving < 30 minutes of vigorous intensity PA ($z = -1.040$, $P = 0.073$) and between 30 to 60 minutes of light/moderate intensity PA ($z = -1.102$, $P = 0.070$). ($z = -1.040$, $P = 0.073$). The MET values of vigorous and light/moderate intensity PA decreased by 48% ($z = -4.334$, $P > 0.001$) and 38% ($z = -5.434$, $P > 0.001$) from before to during the Pandemic. Indoor play,	Not reported.	Fair

No.	Author (Year)	PA Measure; PA Data Reported by ^a	Sample Source (Country)	Sample Size	Age Ranges	Findings	Biological, Environmental, and Social Factors that Affected PA	NIH NHLBI Quality Assessment
						cycling, walking, and at home PA classes reported as types of PA children engaged in during the Pandemic.		
64	Tornaghi, M., et al. (2020).	IPAQ; 1	Italy	1568 children	15-18	Before the Pandemic, 6.8% of children met PA recommendations (60 min of MVPA/day) compared to 14.7% during the Pandemic. From before to during the Pandemic, the percentage of children with a inactive (<700 MET-min/week) (17.8% vs 25.8%) or intense (>2520 MET-min/week) PA level increased (15.8% vs. 19.8%), while the percentage of children with a moderate (700-2519 MET-min/week) PA level decreased (66.3% to 53.6%).	Not reported.	Good
65	Wunsch, K., et al. (2021).	MoMo-PAQ; 1, 2	Germany	1711 children	4-17	Before the Pandemic, male and female children (4-10) met PA recommendations (at least 60 minutes of PA) 4.74 and 4.62 days/week, while during the Pandemic boys and girls met PA recommendations 5.39 and 5.27 days/week. Similarly, the days/week that male and female adolescents (11-17) met PA recommendations increased from 3.90 and 3.55 days/week to 4.08 and 3.96 days/week.	Sedentary time before the Pandemic had a negative association with PA during the Pandemic, independent of age and gender. In contrast, among females and children < 10 years, health related quality of life before the Pandemic was positively associated with PA during the Pandemic.	Good
66	Zorcec, T., et al. (2020).	Questionnaire; 2	Republic of Macedonia	72 parents/ca regivers	7.3±2.89	Among children with cystic fibrosis (CF), asthma, tuberculosis (TB) and allergic rhinitis, the percentage of children who participated in more than 2 hours of PA/day significantly decreased from before to during the pandemic (52.8% and 20.8%, p=0.0001). 11.4% of children before the Pandemic compared to 30.2% of children during the Pandemic participated in only a few hours of PA/week (p=0.0056).	Not reported.	Fair
67	Carroll (2020)	IPAQ; 2	Canada	310 children, 254 families	18 months - 5 years	During lockdown, decreased PA was reported for 52% of children, with an average time spent outdoors and in active play of one hr/day. Parents reported decreased intensity of PA among children during the pandemic.	Parents reported PA was limited during the Pandemic because of a lack of space and variety of tools/toys.	Good
68	Ellis, W. E., et al. (2020).	W. Godin Leisure-Time Exercise questionnaire; 1	Canada	1054 children	14-18	Many participants reported no strenuous (40%) or moderate (21.3%) physical activity over the past week. During the Pandemic, children participated in >5 minutes of mild PA 3.47 times, moderate PA 2.7 times, and strenuous PA 1.93 times.	Not reported.	Good
69	Guerrero, M. D., et al. (2020).	Online survey; 2	Canada	1472 parents	5-17	81.8% of children did not meet the PA recommendation.	Children were more likely to achieve PA recommendations if their parents were able to restrict screen time, had a higher income, or were younger than 43 years old. Furthermore, boys and children with increased outdoor PA and sport during the Pandemic were more likely to achieve PA recommendations.	Good
70	Hemphill, N. M., et al. (2020).	Fitbit (step count)	Canada	109 children	9-16	Children with congenital heart disease obtained 21%-24% less daily steps during early-COVID. Daily steps were consistently below the Canadian PA guidelines of 60	Daily step counts were higher among boys than girls.	Good

No.	Author (Year)	PA Measure; PA Data Reported by ^a	Sample Source (Country)	Sample Size	Age Ranges	Findings	Biological, Environmental, and Social Factors that Affected PA	NIH NHLBI Quality Assessment
						minutes of MVPA/day (approximately equivalent to 12,000 steps/day).		
71	McCormack, G. R., et al. 2 (2020).	Online questionnaire;	Canada	328 parents	5-17	During the Pandemic, children met PA recommendations (≥ 60 min of MVPA) an average of 3.48 ± 2.41 days/week. Children played 4.52 ± 2.75 days in the past week and went to a park on 4.85 ± 6.39 days in the past month. Meanwhile, 18.3% of children reported 0 days of MVPA/week, 16.5% had 0 days of play per week, and 34.5% had no visits to a park in the past month. 48.8% of parents reported their child's PA at home increased and 32.9% of parents reported that their child's PA was the same compared to before the Pandemic. 38.7% of parents reported their child's PA outdoors increased, while 39% of parents reported their child's PA outdoors decreased. 52.7% of parents reported their child's play at a park decreased, while 15.5% of parents reported their child's play at a park increased. 53.7% of parents reported their child's play in public spaces decreased, while 9.5% of parents reported their child's play in public spaces increased.	Children of parents with high COVID-19 anxiety had fewer days at the park and less PA both in the home (14.2% vs. 25.6%, $p < .05$) and outdoors (33.2% vs. 49.6%) compared to children of parents with low COVID-19 anxiety (3.00 days vs. 5.42, $p < .05$).	Fair
72	Mitra, R., et al. 2 (2020).	Online survey;	Canada	1472 parents	5-17	More youth than children experienced a decrease in physical activity-related movements during the pandemic, including walking/biking, outdoor or indoor physical exercise and outdoor play. The majority of children reported a decrease in walking or biking (53.2%), PA or sport outside (63.8%), and outdoor play (51.2%) during the Pandemic. In contrast, the majority of children reported PA or sport inside stayed the same (40.5%), while indoor play increased (53.1%).	Compared to children who reported decreased outdoor activity during the Pandemic, a greater proportion of children who reported increased outdoor activity met PA guidelines (11.6% vs 26.5%, $p < 0.001$) (60 minutes of MVPA/day). Younger children, children with a higher household income, children living in houses (vs apartments), living in multi-child households, and living further from major roads were more likely to cluster into the increased outdoor activity group. Gender was not associated with changes in outdoor activity. While neighborhood density was negatively associated with clustering into the increased outdoor activity group, access to a park within 1 km increased the likelihood of children living in high density neighborhoods clustering into the increased outdoor activity group (OR = 1.19).	Good
73	Moore, S., et al. 2 (2020).	Online survey;	Canada	1472 parents	5-17	During the Pandemic, 23.8% of children (5-13 years) and 13.2% of youth (14-17 years) met PA recommendations (60 minutes of MVPA/day). Children and youth attained 60 minutes of MVPA an average of 3.55 and 2.59 days/week, respectively. PA included activities such as walks or bikes in the neighborhood, PA or sport outside and inside, and outdoor play. Overall, children and youth had decreased PA and outside time during the Pandemic, with the greatest reduction in outdoor PA and sport. In contrast, children and youth reported increased indoor play. Among outdoor activities, biking (6.1%), walking/hiking (5.5%), and sports (3.5%) increased the most during the pandemic.	A greater proportion of children (23.8%) met PA recommendations than youth (13.2%). More boys (5-11 years) met PA recommendations than girls (27.9% boys, 19.0% girls). Younger parental age, parental encouragement, parental co-participation in PA, parental cohabitation, living in a detached house (vs. apartment) and having a dog were positively associated with PA.	Good

No.	Author (Year)	PA Measure; PA Data Reported by ^a	Sample Source (Country)	Sample Size	Age Ranges	Findings	Biological, Environmental, and Social Factors that Affected PA	NIH NHLBI Quality Assessment
74	Bazett-Jones (2020)	Online questionnaire; 1	US	287 children	9-19	During COVID-19 restrictions, runners reported shorter distance runs ($p < 0.001$), fewer runs ($p < 0.001$), and fewer intense runs ($p < 0.001$) per week.	Not reported.	Fair
75	Dunton, G. F., et al. (2020).	Online survey; 2	US	211 children	5-13	During the Pandemic, the most reported forms of PA were free play/unstructured activity (i.e. running around, tag) (90% of children) and walking (55% of children). Remote/streaming services were also used for PA programs (i.e. martial arts, dance, yoga classes) (28.9% of children), with older children (ages 9-13) five times more likely to participate in virtual team sports training sessions compared to younger children (ages 5-8)(OR = 5.40, 95% CI [1.70,17.15], Wald = 8.19, $p = .004$)Wald = .288, $p = .633$). Younger children (age 5–8) were more likely to participate in free play/unstructured physical activity, biking, and scootering/skateboarding/roller skating, while older children (ages 9–13) were more likely to participate in circuit training/conditioning. During the Pandemic, the likelihood of participating in PA at home (OR = 2.49, 95% CI[1.35, 4.60], Wald = 8.593, $p = .003$ and in the neighborhood (i.e. on sidewalks and roads)(OR = 1.92, 95% CI [1.04,4.60], Wald = 4.28, $p = .038$) increased compared to before the Pandemic. In contrast, the likelihood of participating in PA at parks and trails decreased (OR = 0.47, 95% CI [0.23, 0.97], Wald = 4.22, $p = .040$).	Parents perceived greater decreases in PA among older children (ages 9–13) compared to younger children (ages 5–8).	Good
76	Garcia, J. M., et al. (2020).	Online survey; 1	US	9 children	14-19	Prior to the pandemic, participants reported more days that they participated in 60 minutes or more of PA more frequently (4.17 days vs. 2.27 days; $p = 0.0006$) and participated in more activities (3.4 activities vs 2.11 activities; $p = 0.007$) than during the pandemic. 78% of the participants felt that their PA decreased.	The participants with decreased PA reported school PA program cancellations and fear of going outdoors as barriers to PA. The participant with increased PA reported it was because his family started riding bikes together.	Good
77	McGuine, T. A., et al. (2020).	Pediatric Functional Activity Brief Scale (PFABS); 1	US	13002 children	3-19	Among team sport athletes, lower levels of PA were reported compared to athletes involved in individual sports.	Lower levels of PA were reported among girls, athletes in the 12th grade, athletes who participated in team sports, and athletes from more impoverished areas.	Good
78	Pavlovic, A., et al. (2021).	Online questionnaire; 3	US	2440; 1789 PE teachers, 64 district administrators, 62 school administrators, and 3 nurses	not specified (grades 3-12)	During the Pandemic, 79% of participants reported students PA was "significantly less" or "somewhat less". Frequently reported online PA resources included YouTube (29%),virtual learning platforms (27.9%), and PE teacher-led virtual classes (25.6%). Among schools that closed due to the Pandemic, barriers to PA included student access to virtual learning and communication between teachers and students. Among schools that remained open, barriers to PA were social distancing guidelines and access to equipment and space for PA.	There were no significant differences in PA between children in different education levels (i.e., elementary, middle, or high schools) or regions of the U.S.	Good

No.	Author (Year)	PA Measure; PA Data Reported by ^a	Sample Source (Country)	Sample Size	Age Ranges	Findings	Biological, Environmental, and Social Factors that Affected PA	NIH NHLBI Quality Assessment
79	Tulchin-Francis, K., et al. (2021).	Weighted Leisure-Time Score Index (unvalidated modified Godin Leisure-Time Exercise Questionnaire (mGodin); 2	US	1083 parents	3-18	While light PA was unchanged, MVPA of children significantly decreased during the Pandemic (before: 46.7, during: 34.7, $p < 0.001$). For all ages, PA with friends, yard and neighborhood play (18.1%–39.8%, $p < 0.05$), and indoor child and parent-led play (6.8%–23.4%, $p < 0.05$) decreased. Preschool and elementary children reported decreased PA with family (8.3%–23.5%, $p < 0.05$) and a 7.1% to 20.1% increase in independent, virtual PA. Among preschool and elementary girls (20.3% and 5.7%, $p < 0.05$), as well as, elementary and middle school boys (5.1% and 13.8%, $p < 0.05$), increased yard play was reported. In contrast, there were no significant increases in outdoor PA among high school students. Before the Pandemic, 78.6% of children participated in organized sports, while during the Pandemic only 10.6% participated in organized sports.	The Pandemic had the lowest impact on PA (based on quantity, variety, and intensity of PA) of preschool students and the highest impact on PA of high school students. Boys had higher reductions in PA than girls.	Good
80	de Matos (2020)	IPAQ; 1	Brazil	69 children	8-18	Weekly energy expenditure was significantly reduced among children (male: ES: 3.02, CI95%: [1.93; 4.12], $p < 0.0001$; female: ES: 3.26, CI95%: [1.82; 4.69], $p < 0.0001$) and adolescents (male: ES: 4.25, CI 95%: [3.06; 5.44], $p < 0.0001$; female: ES: 4.01, CI 95%: [3.02; 5.00]; $p < 0.0001$) during the Pandemic.	Not reported.	Fair
81	Sa, C., et al. (2020).	Online survey; 2	Brazil	816 children	0-12	Most parents reported a decrease in PA during the Pandemic, with children doing either much less PA (46.1% of parents) or less PA than during a normal school year (37% of parents). Children participated in 0.4 ± 0.6 to 0.7 ± 1.0 hours of PA/day during the Pandemic.	Older children had lower levels of playing with PA than younger children. There were no significant relationships between age and sex and organized or outdoor PA.	Good
82	Siegle, C. B. H., et al. (2020).	Online questionnaire; 2	Brazil	816 children	0-12	During the Pandemic, the highest average percentage of daily PA was $10.3 \pm 16.2\%$ among boys ages 0 to 2, while the lowest average percentage of daily PA was 3.9 ± 5.7 among girls ages 6 to 9. (%PA = playtime w/ PA + PA (planned, inside/outside the house, dog walking) / (playtime w/ PA + PA + playtime w/o PA + playtime on screens + intellectual (school))).	Percentage of PA (%PA) was higher among children with large external spaces at home than children with small external space ($p < 0.001$) or with no external space ($p < 0.001$). Older children had lower %PA than younger children. There was no significant effect of gender on differences in %PA.	Good
83	Aguilar-Farias (2020)	Online survey; 2	Chile	3157 children	1-5	For all age groups, the average duration of PA decreased by 0.75 hrs/day (-0.75 [CI 95% $-0.81, -0.70$] h/day).	Greater reductions in PA were reported for children with a more educated main caregiver or who lived in an apartment. Smaller reductions in PA were reported for children living with more children or with five or more people, children in rural areas, and children with space at home to play. While there was no effect of gender on PA, older children had less PA.	Good
84	Reece, L. J., et al. (2020).	Online survey; 2	Australia	16177 parents	4-18	During the Pandemic, 31% of participants reported children's PA decreased a lot, while 39% reported children's PA decreased a little. For PA, 82% of children were active at home,	Not reported.	Fair

No.	Author (Year)	PA Measure; PA Data Reported by ^a	Sample Source (Country)	Sample Size	Age Ranges	Findings	Biological, Environmental, and Social Factors that Affected PA	NIH NHLBI Quality Assessment
						while 52% of children were active in public spaces.		
85	Cahal (2021)	Online questionnaire; 2	Israel	445 children	0-18	PA decreased in 53.9% of patients with chronic respiratory disorders (i.e. asthma, recurrent pneumonia).	Older patients (>5 years old) had decreased PA compared to younger patients (p<.001).	Fair
86	Chaturvedi (2021)	Online survey; 1	India	303 children	7-17	Children spent an average of 0.82 hrs/day on fitness during the Pandemic.	Not reported.	Good
87	Ghanama h, R. and H. Eghbaria-Ghanama h (2021).	Online survey; 2	Israel	382 parents	5-11	Children spent less time in physical activities during the Pandemic than before. (before: 2.88 hours/day during: 2.02 hours/day, p<0.001)	Children spent less time in PA when a family member was diagnosed with COVID-19.	Good
88	Masi, A., et al. (2021).	Online survey; 2	Australia	302 caregivers	2-17	68% of caregivers for children with neurodevelopmental disabilities reported a decrease in exercise.	Not reported.	Fair
89	Munasin ghe, S., et al. (2020).	PACE +, smartphone sensors (pedometer, MBAR); 1	Australia	582 children	13-19	PA decreased from before to during the Pandemic (odds ratio [OR] = .53, 95% confidence interval [CI] = .34–.83), with a concurrent decrease in the average number of steps/day.	Not reported.	Fair
90	Nathan, A., et al. (2021).	Online survey; 2	Australia	157 children	5-9	Compared to before the Pandemic, the total minutes of PA/week did not significantly change during the Pandemic. However, there was a 65.7% and 65% reduction in the duration and frequency of organized PA (-124.6 min/week, p<0.001; -1.3 times/week, p<0.001). In contrast, the duration and frequency of total (23.1% or 146.3 min/week, p=0.005 and 30% or 1.8 times/week, p<0.001) and home-based (58.7% or 201 min/week, p<0.001) unstructured PA increased. Other activities that significantly increased from before to during the Pandemic included outdoor play (i.e., in the yard or street around the house, in a park or outdoor recreation area) and home indoor play. There was a 95% increase in the min/week reported for outdoor play in a park, playground, or outdoor recreation area.	Parents reported facilitators of PA during the Pandemic as increased time at home to support PA, more children outside playing together, more utilization of local PA resources (i.e., parks), more unstructured PA, and more parental encouragement.	Good
91	Parker, K., et al. (2021).	Online survey; 1	Australia	963 children	13-17	During the Pandemic, 7.2% of children met PA recommendations (60 min of MVPA/day). Among the 26.5% of children who used digital platforms for PA, streaming services were used by 40.0%, online classes were used by 30.2%, and subscriber fitness programs were used by 13.7% of those children. The median frequency and duration of digital platform use for PA were 4 (IQR 3-7) times per week and 120 (IQR 60-260) min/week.	Children who used digital platforms for PA during the Pandemic were over 2 times more likely to meet PA recommendations.	Good
92	Sciberras, E., et al. (2020).	Online survey; 2	Australia	213 children	5-17	Among kids with ADHD, there was decreased exercise (Odds Ratio (OR) = 0.4; 95% CI 0.3–0.6) and outdoor	Not reported.	Good

No.	Author (Year)	PA Measure; PA Data Reported by ^a	Sample Source (Country)	Sample Size	Age Ranges	Findings	Biological, Environmental, and Social Factors that Affected PA	NIH NHLBI Quality Assessment
						time (OR = 0.4; 95% 0.3–0.6) compared to pre-COVID.		
93	Zhao, Y., et al. (2020).	Online questionnaire; 1, 2	China	738 students, 1062 parents	11.0±1.7	During the Pandemic, 26% of children reported 0 hours, 18% reported <1 hour, 37.9% reported 1–2 hours, and 18% reported more than 2 hours of daily outdoor activity. In contrast, 43.7% of parents reported their child's daily outdoor activity was 0 hours. 42.9% of children in grades 7–9 reported 0 hours of daily outdoor activity compared to 24.1% of children in grades 1–3 and 23.8% of children in grades 4–6.	Not reported.	Fair
95	Wu, X., et al. (2021).	Telephone questionnaire; 2	China	43 children	0–18	Among children with type 1 diabetes, 44.1% reported PA decreased during the Pandemic. Before the Pandemic, outdoor activities (i.e., cycling, basketball) were predominant forms of exercise. However, during the Pandemic, the predominant form of exercise was indoor activities (i.e., pacing, skipping rope).	Not reported.	Fair
96	Zhang, X., et al. (2020).	IPAQ-SF; 1	China	9979 children	9–14	Students obtained an average of 23.19 minutes of moderate-to-vigorous PA (MVPA)/ day (1193.02 ± 1621.88 MET-min/week). Vigorous PA made up 42.74% (510.40 ± 934.18 MET-min/week) of total PA, while walking for PA made up 24.19% (288.60 ± 613.08 MET-min/week).	Girls participated in significantly more MVPA and moderate PA than boys during the Pandemic (p<0.01). Similarly, younger children in grade 4 participated in significantly less MVPA and moderate PA than older children in grades 5 and 6. (p<0.01)	Good
97	Abid (2021)	Ricci and Gagnon sedentary behavior questionnaire; 1	Tunisia	100 children	5–12	Compared to before the Pandemic, sports and recreational PA decreased by 35%, daily PA decreased by 16% for boys and 27% for girls, and total PA decreased by 7% for boys and 17% for girls during the Pandemic (p<0.001).	There was no significant effect of gender on PA.	Fair
98	Shinomiy a, Y., et al. (2021).	Online survey; 2	Japan	2019: 2017 children, 2020: 295 children	18–30 months	The percentage of leisure time that consisted of outdoor play decreased from an average of 64.0 ± 26.0% before the Pandemic (March 2019) to 61.0±27.1% during the Pandemic (March 2020) (p=0.058).	Children who stayed at home had less outdoor play than children who received childcare at a nursery school (p=0.019).	Good
99	Xiang, M., et al. (2020).	Global Physical Activity questionnaire (GPAQ); 1	China	2426 children	6–17	Before the Pandemic, PA decreased from a median of 540 min/week to 105 min/week during the Pandemic (-435 min/week, p<0.001). The percentage of children who met PA recommendations (at least 60 min of PA/day) decreased from 60% to 17.7% during the Pandemic (-42.3%).	Not reported.	Fair
100	Yang, S., et al. (2020).	IPAQ-long form; 1	China	2824 children	17.5 ± 1.2	High school students reported their MVPA (80.4%), active transport for commuting and errands (77.4%), housework activity (50.1%), and walking for leisure (79.9%) stayed the same. The hrs/week that high school students participated in MVPA remained	Not reported.	Fair

No.	Author (Year)	PA Measure; PA Data Reported by ^a	Sample Source (Country)	Sample Size	Age Ranges	Findings	Biological, Environmental, and Social Factors that Affected PA	NIH NHLBI Quality Assessment
						unchanged at an average of 1.5 hrs/week.		
101	Elnaggar, R. K., et al. (2020).	PAQ-A; 1	Saudi Arabia	63 children	14-18	From baseline to follow-up, the PA level (PAL) of all participants was significantly reduced (BL-PAL: 3.05 ± 0.54; FL-PAL: 2.77 ± 0.47; P < .001).	While PAL decreased for both genders, only the PAL reduction was significant in boys (BL-PAL: 3.20 ± 0.57; FU-PAL: 2.76 ± 0.49; P < .001). At baseline, boys achieved significantly higher PAL than girls (P = .014), but at follow-up both genders had similar PAL (P = .86).	Fair
102	Jia, P., et al. (2021).	IPAQ-long form; 1	China	2824 children	16-18	Compared to before the Pandemic, the frequency of active transport (i.e., commuting and errands) among high school students decreased from 1.3 to 0.9 days/week (p<0.001), while leisure-time walking decreased from 1.0 to 0.7 days/week (p<0.001) during the Pandemic. Similarly, the frequency of moderate-to-vigorous housework decreased from 2.3 to 1.9 days/week (p < 0.05) and leisure-time moderate-to-vigorous PA decreased from 0.7 to 0.65 days/week (p < 0.001).	Not reported.	Good

Legend:^a 1 = child reported measure; 2 = parent reported measure, 3 = other secondary subjective reported measure. Abbreviations: PA, physical activity; NIH NHLBI, National Institute of Health National Heart, Lung, and Blood Institute; CI, confidence interval; h/day, hrs/day; MVPA, moderate-to-vigorous physical activity; M, mean; SD, standard deviation; PAL, physical activity level; ES, effect size; min/d, min/day; OR, odds ratio; Wald, Wald test; BL, baseline; FU, follow up; P.E., physical education; min/wk, min/week; ml.kg⁻¹.min⁻¹, millimeters per kilogram per minute; h/d, hrs/day; km, kilometers; ADHD, attention-deficit/hyperactivity disorder; LPA, light physical activity; MET, metabolic equivalent task.

The sample sizes ranged from 9 children⁷⁶-16177 parents.⁸⁴ Most articles (n=32)^{33-35,38-41,43,47,48,50,51,58,60,62,63,67,70,71,74,75,81,82,85-93} had a sample size between 100-1000, but 2 articles had a sample size >10000.^{77,84} As the present study is focused on children, most of the articles are inclusive of age groups between 0-18 years. However, 5 articles included participants ≤19 years because the articles specified their sample as either adolescents (n=4)^{58,76,77,89} or school-aged youth (n=1).⁷⁴ Because adolescents are defined by the WHO as individuals 10-19 years old,⁹⁴ these articles were included despite the exclusion criteria of ≤18 years. One article did not specify the age of participants,⁷⁸ however, specified participants were in grades 3-12, and was ultimately included.

Most participants were children without chronic health conditions, but some articles focused on characterizing PA during the pandemic for children with various health conditions (e.g., autism spectrum disorder, congenital heart disease, chronic respiratory disorders, obesity, type 1 diabetes, attention-deficit/hyperactivity disorder; n=11).^{52,54,56,63,66,70,76,85,88,92,95} Some articles examined PA during the pandemic only (n=13),^{35,45,55,57,61,68,69,77,82,86,91,93,96} while others compared PA from before to during the pandemic (n=56).^{33,34,36-44,46-54,56,58-60,62-67,70-76,78-81,83-85,87-90,92,95,97-102}

PA data was reported by either the child (n=28),^{38-41,43,48,52,56,58-62,64,68,74,76,77,80,86,89,91,96,97,99-102} the parent or guardian

(n=32),^{34,35,37,42,44,47,49-51,53-55,63,66,67,69,71-73,75,79,81-85,87,88,90,92,95,98} a different secondary subjective individual such as a school administrator (n=1),⁷⁸ or a combination of these groups (n=5).^{36,45,57,65,93} Based on the NIH NHLBI Study Quality Assessment Toolbox,³² 50 articles received a rating of "good," while 19 articles received a rating of "fair" [Table 1](#).

PA Measures

PA measures used in the included articles are depicted in [Table 2](#). Online surveys were predominantly used to collect PA data. Common PA questionnaires included the International Physical Activity Questionnaire (IPAQ, short or long form; n=8)^{58,63,64,67,80,96,100,102} and the Physical Activity Questionnaire for Adolescents (PAQ-A; n=5)^{38-41,101} or for children (PAQ-C; n=2).^{44,45} The PAQ-A designates a physical activity level (PAL) based off a scale of 1-5: 1 designating no activity or a low activity level and 5 designating a high activity level. A PAL score below 2.73 suggests an insufficient PAL and a score above 2.73 suggests a sufficient PAL. A few surveys took place in-person (n=2)^{43,66} or over the phone (n=4).^{37,54,56,95} Seven studies used objective measures to collect PA data, including the use of a Fitbit,⁷⁰ accelerometer,^{33,37,48,62} smartphone sensors (e.g., pedometer),⁸⁹ or a 20-meter shuttle run test to measure VO2 max.⁴⁶ The most prevalent FITT principle collected to determine PA was time (e.g., duration of PA; n=46),^{33,36,42,45,47,48,50,52-68,71-73,75,76,80-84,86,87,89-93,96-100} followed by frequency (n=30),^{38,39,43-45,49,51,52,59,60,63-65,68,69,71,73,74,76,77,79-81,84,90-92,96,102}

type of exercise (n=30),^{36,38,39,43-45,51,59,60,62,63,68,71-73,75,77-79,84,89-92,95-98,100,102} and intensity (n=25).^{33,36,45,48,52,53,59,62-64,67,68,71-74,79,80,84,91,96,97,99,100,102}

Types of PA

Among the articles included, there were conflicting results concerning the effects of the pandemic on the types of PA in which children participated. Five articles reported that outdoor PA declined during the pandemic,^{71-73,95,98} while 3 studies reported that outdoor PA increased.^{37,51,90} The types of outdoor PA included activities such as walking, biking, outdoor play, playing at parks or other public spaces, and sports. Eight studies reported that many children (e.g., 82%)⁸⁴ were active at home.^{60,61,63,71,75,84,90,95} Some studies reported the use of online platforms such as streaming services (e.g., YouTube), virtual classes (e.g., yoga),^{45,75} or other virtual PA forms⁷⁹ to accumulate PA during the pandemic.^{78,91} In a US study, the most common forms of PA during the pandemic were unstructured play (90%) and walking (55%).⁷⁵ In contrast, a Chinese study reported the most common form of PA changed from outdoor activities (e.g., cycling) to indoor activities (e.g., rope skipping).⁹⁵

Extent to Which PA Changed During the Pandemic

Several studies (n=10) reported most children obtained <30 min/day or <3 hrs/week of PA,^{33,35,42,47,56,61,66,68,96,99} with 2 reporting most children participated in PA 0 days/week.^{35,68,99} Of the 12 articles^{37,45,49,50,53,57,62,64,69,73,91,99} that reported the percentage of children meeting PA guidelines (i.e., 60 MVPA min/day), 1 article reported <10%,⁹¹ 9 articles reported 10-30%,^{45,49,53,57,62,64,69,73,99} and 2 articles reported >50%^{37,50} of children met PA recommendations during the pandemic. All articles that utilized the PAQ-A to assess PA (n=5),^{38-41,101} reported a significant decrease in PAL.

Of the articles that compared PA before the pandemic to during the pandemic (n=56),^{33,34,36-44,46-54,56,58-60,62-67,70-76,78-81,83-85,87-90,92,95,97-102} 89% reported a decrease in PA (i.e., frequency, intensity, type/duration, type) of children. Among these articles, the proportion of participants that reported a decrease in PA behaviors during the pandemic ranged from 42%⁴³- 95%.⁴⁸ One study reported that although the total minutes of PA/week did not significantly change, the duration and frequency of organized PA decreased, while unstructured PA and outdoor play increased.⁹⁰

Similarly, another study found 80.4% of high school students reported their MVPA stayed the same at 1.5 hrs/week.¹⁰⁰ Two studies^{37,59} reported an increase in PA (e.g., duration, frequency of meeting PA recommendations/week) of children (n=4), 1 of which reported a specific increase in outdoor time compared to before the pandemic.³⁷ One study reported 90.3% of preschoolers (3- 5 years old) met PA guidelines of ≤180 minutes of PA/day, with 60 minutes of MVPA/day.³⁷ The articles that reported PA (e.g., duration, frequency of meeting PA recommendations per week) increased during the pandemic had samples from Sweden,³⁷ Canada,⁷¹ and Germany.^{59,65}

Conversely, four studies from Canada,^{67,70,72,73} four studies from Australia,^{84,88,89,92} and one study from Germany⁴³ reported PA (e.g., frequency, intensity, daily steps, meeting PA recommendations) decreased. For studies that included children with pre-existing health conditions (n=11),^{52,54,56,63,66,70,76,85,88,92,95} PA (e.g., daily steps, duration, frequency) decreased compared to before the pandemic. For example, among children with obesity in Italy, sports participation decreased by 2.30±4.60 hrs/week (p=0.003).⁵⁴


Biological, Environmental, and Social Factors Affecting PA

Common trends were evident among the studies of this review that related factors such as gender, age, and the environment outside as well as inside the home to the PA of children during the pandemic. In general, boys had higher levels of PA (e.g., more minutes of PA/day or more likely to meet PA recommendations) than girls. A study in Tunisia found that total PA (score from the Ricci and Gagnon sedentary behavior questionnaire based on duration and intensity of PA) decreased by 17% and 7% among girls and boys, respectively.⁹⁷ Older children tended to have greater reductions in time/duration of PA compared to younger children. For example, in 2 Spanish studies, PA decreased an average of 43.3 min/day among preschoolers aged 4- 6 years old,³³ while PA decreased an average of 91 ± 55 min/day among older children aged 8- 16 years old.⁴⁸

There were mixed results for PA of children and parents' socioeconomic status (e.g., lower levels of PA were reported almost equally for both high and low parental income and education). Children who lived in a rural areas⁸³ or had more outdoor space to play^{36,48,55,82} had lower reductions in PA (e.g., duration, less likely to report decreased PA, greater percentage of day spent participating in PA). For example, children with access to an outdoor area at home had a 1.4 ± 0.1 hrs/day reduction in PA compared to a 1.7 ± 0.1 hrs/day reduction in children without access (p<.01).⁴⁸ In contrast, children who lived in a high-density neighborhood⁷² or urban area^{36,41} had greater reductions in PA (e.g., more likely to report decreased PA, less likely to meet PA recommendations, greater reduction in PAL). Children who lived in an apartment had lower levels of PA (e.g., decrease in duration of PA or less likely to meet PA recommendations) compared to children who lived in a house.^{72,83}

The environment inside the home also seemed to affect PA of children during the pandemic. One study reported that children who lived in a home with more people had greater reductions in duration of PA,⁸³ while 3 studies reported children who lived in multi-child households (e.g., with siblings) participated in more PA (e.g., greater duration or greater percentage of day spent participating in PA)^{55,83} or more frequent outdoor play.^{51,72} Children who had less family conflict, more parental support, and more family engagement in PA had greater durations of PA.^{53,73,76} However, children who had more conflict with their parents or who had parents with higher stress levels had less PA (e.g., shorter duration of PA or less likely to achieve a sufficient PAL).^{38,53}

Figure 2. Sample Source (Countries) of Included Articles.



Discussion

The present study aimed to address the types of PA children participated in, the extent to which PA (e.g., frequency, intensity, duration, and type) changed, and factors affecting PA during the pandemic. Schools, active transportation (e.g., walking to and from school), afterschool programs, and sport programs were predominant sources of MVPA prior to the pandemic.^{103,104} The stay-at-home orders and social distancing policies put in place to mitigate the spread of COVID-19 restricted the ability of children to obtain PA through these previously popular sources. While the studies in this review revealed walking, outdoor activity, and activity at home were types of PA children engaged in during the pandemic, a study before the pandemic found that children were less likely to participate in routine PA (e.g., going to the gym, walking) compared to unstructured play.⁵ Social distancing restrictions could have contributed to shorter durations of PA and overall decreased participation, despite walking becoming a common source of PA during the pandemic.⁷⁵ In addition, some studies reported many children were active at home (e.g., indoors or not at a park/public space).^{60,61,63,71,75,84,90,95} Studies conducted prior to the pandemic found that indoor PA (e.g., at home, at school) tends to be a greater source of lighter intensity PA,¹⁰³ which could help to explain the reduction in children able to meet MVPA recommendations during the pandemic.

The results of this review suggested that while there was heterogeneity in the extent to which the pandemic affected PA (e.g., frequency, intensity, duration, and type) in children, overall, there was a decrease in PA of children globally. Although the impact of the pandemic on PA of children is still evolving, trends before the pandemic have shown that children tend to have lower levels of PA when they are not in school.^{13,14} Studies that investigated changes in MVPA during time out of school (e.g., weekends or a 3-week school break) found that MVPA decreased by 10-14 minutes compared to week days during the school year.^{13,105} In contrast, the reductions in PA presented by the studies in this review ranged from 32-91 min/day.^{33,48,54,83,99} These results revealed that the decreases in time/duration of PA during the pandemic may have been greater than those typically observed during days when children are not in school. Prior to the pandemic, global PA levels of children were already low, with 81% of children insufficiently physically active.¹² Of the studies that

Table 2 Characteristics of the Physical Activity Data for the Included Articles (n=69).

Study Characteristic	Total Articles	Articles
PA Data Reported by:		
Child	28	38-41, 43, 48, 52, 56, 58-62, 64, 68, 74, 76, 77, 80, 86, 89, 91, 96, 97, 99-102
Parent	32	34, 35, 37, 42, 44, 47, 49-51, 53-55, 63, 66, 67, 69, 71-73, 75, 79, 81-85, 87, 88, 90, 92, 95, 98
Child and Parent	5	36, 45, 57, 65, 93
Other (Fitbit, accelerometer, VO2, smartphone sensor)	7	33, 37, 46, 48, 62, 70, 89
Teacher and Parent	1	58
Teacher and School Administrator	1	78
PA Measure		
IPAQ (short or long form)	8	59, 64, 65, 68, 81, 97, 101, 103
PAQ-A	5	39-42, 102
PAQ-C	2	44, 45
PACE+	2	52, 89
Godin Leisure-Time Exercise Questionnaire (or modified version)	2	68, 79
Other online survey	39	34-36, 42, 47-51, 53, 55, 57, 60-62, 69, 71-78, 81-88, 90-93, 97-99
MoMo-PAQ	2	59, 65
Telephone interview	4	37, 54, 56, 95
In-person questionnaire	2	43, 66
Accelerometer (GENEActiv tri-axial, ActiGraph GT3X)	4	34, 38, 49, 63
FitBit	1	70
Smartphone sensors (pedometer, MBAR)	1	89
VO2 max (20 meter shuttle run test)	1	46
Physical Activity Time Frame		
During the COVID-19 pandemic	13	35, 45, 55, 57, 61, 68, 69, 77, 82, 86, 91, 93, 96
Compared PA before to during the COVID-19 pandemic	56	33, 34, 36-44, 46-54, 56, 58-60, 62-67, 70-76, 78-81, 83-85, 87-90, 92, 95, 97-102
Did PA increase or decrease?		
PA decreased	50	33, 34, 36, 38-44, 46-54, 56, 58, 60, 62-64, 66, 67, 70, 72-76, 78-81, 83-85, 87-89, 92, 95, 97-102
PA increased	4	37, 59, 65, 71
PA stayed the same	2	90, 100
FITT Principle reported		
Frequency	30	38, 39, 43-45, 49, 51, 52, 59, 60, 63-65, 68, 69, 71, 73, 74, 76, 77, 79-81, 84, 90-92, 96, 102
Intensity	25	34, 37, 46, 49, 53, 54, 60, 63-65, 68, 69, 72-75, 80, 81, 85, 92, 97, 98, 100, 101, 103
Time	46	33, 36, 42, 45, 47, 48, 50, 52-68, 71-73, 75, 76, 80-84, 86, 87, 89-93, 96-100
Type	30	37, 39, 40, 44-46, 52, 60, 61, 63, 64, 69, 72-74, 76, 78-80, 85, 90-93, 96-99, 101, 103

Legend: Abbreviations: PA, physical activity; IPAQ, International Physical Activity Questionnaire; PAQ-A, Physical Activity Questionnaire for Adolescents; PAQ-C, Physical Activity Questionnaire for Children; PACE+, Patient-centered Assessment and Counseling for Exercise plus Nutrition; MoMo-PAQ, Motorik-Modul Physical Activity Questionnaire.

reported a decrease in PA during the pandemic, the percentage of children who did not meet PA recommendations ranged from 80%- 99.7%.^{45,49,62,69,73,91} This suggests that global levels of PA in children may have worsened during the pandemic.

The results of the current study indicate that during the pandemic, boys, and younger children tended to have higher levels of PA (e.g., more likely to meet PA recommendations), which is consistent with PA trends that existed prior to the pandemic.^{104,106-109} Previous studies have found that self-efficacy is an important predictor of PA.^{108,110} Not only are girls less likely to participate in PA,¹¹¹ but they also tend to have a lower perceived ability to overcome PA barriers.¹¹² The social cognitive theory suggests that self-efficacy moderates the relationship between the social and physical environment and changes in PA, such that individuals with a high self-efficacy for overcoming PA barriers should be more likely to participate PA despite changes in the environment.¹¹³ With the changes in the social and physical environment surrounding children during the pandemic, self-efficacy for overcoming PA barriers may have played a role in the observed changes in PA. Similarly, lower levels of self-efficacy in girls may also be due to lower levels of social support to participate in PA compared to boys.¹¹¹

The relationship between self-efficacy, social support, and PA may also apply to age-related changes in PA. Several studies prior to the pandemic reported that children and adolescents with higher levels of peer and family support had higher levels of PA (e.g., frequency, more likely to meet PA recommendations).^{5,114-116} For example, one study found that adolescents with higher parental support were 40% more likely to achieve sufficient PA (60 minutes of MVPA/day).¹¹⁵ While it has been established that PA (e.g., proportion of children meeting PA recommendations) decreases with age,¹¹⁷⁻¹²⁰ 1 such study reported that adolescents with greater self-efficacy had lower age-related reductions in PA.¹¹⁸ The benefits of social support on child PA behaviors could have been especially critical in providing additional support to children navigating barriers to PA during the pandemic. In addition, a past study also found that older girls prefer to do PA at school or in a community setting than at home and participate in team sports.¹²¹

The changes in PA behaviors during the pandemic were likely due to the intersectionality of many social factors that resulted in decreased PA (e.g., frequency, intensity, duration), especially among girls and older children. Greater participation in PA among children with greater social support could also explain why children living in multi-child home had higher levels of PA (e.g., frequency, duration) during the pandemic. Studies before the pandemic have shown that children with siblings or other children to play with participated in more spontaneous and unstructured play.¹²²⁻¹²⁴ More children inside the home may have been a protective factor for PA in children during the pandemic, as it could have offered greater opportunities and incitement to participate in PA.

The studies in this review suggested that during the pandemic, children in rural areas had lower reductions in PA (e.g., duration)

compared to children in urban areas. While some studies before the pandemic have shown MVPA of children was higher in environments with less urbanization,^{125,126} one study suggested that children in suburban or rural areas have less PA.¹¹⁰ The inconclusive relationship between rurality and PA in children could be due to the lack of a global definition of rurality and different distributions of built environments that are conducive to PA in different countries. A study of low- and medium-income countries in Africa found that urban development led to changes in the environment that were less favorable for outdoor PA.¹²⁷ While outdoor time has been shown to be positively related to PA,¹²⁸⁻¹³⁰ rural environments do not guarantee higher levels of outdoor time. Instead, rural environments may also create barriers to PA such as limited quantity of and access to resources that promote PA, such as playgrounds, parks, trails, and other recreational areas. Another study found that differences in MVPA between children in rural and non-rural areas of the U.S. were mediated by differences in neighborhood resources.¹³¹ Rather than comparing PA levels by rurality, more general characteristics of the PA environment such as quantity of and accessibility to community recreational areas, neighborhood safety, traffic levels, and walkability may be more helpful in identifying global factors that affected PA in children during the pandemic.

In conclusion, the purpose of the present review was to synthesize the global impact of the pandemic on PA of children. This review adds to the current body of literature on the effects of the pandemic on PA by specifying changes in the frequency, intensity, duration, and type of PA performed by children, as well as the proportion of children who met PA recommendations. The greatest strength of this systematic review is the range of PA data presented for a wide geographic distribution. Information about different changes in PA (e.g., frequency, intensity, duration, type), as well as characterizations of PA during the pandemic may help to create a succinct picture of the impact of the pandemic on PA in children. A limitation of the systematic review process is some relevant studies may have been missed in the database search due to database selection and exclusion of studies not published in English. In addition, the use of online surveys, the most frequent data-collecting method used in the included studies, is an important limitation of this review, as surveys can only report subjective measures of PA. Furthermore, studies that asked participants to recall PA behaviors prior to the pandemic could be inaccurate. Moreover, online surveys may not have been accessible to individuals of low socioeconomic status who may not have access to the internet. Limited representation of individuals of low socioeconomic status could, therefore, affect the generalizability of the results of this review.

While many PA trends that existed prior to the pandemic persisted, the further reductions in already low global PA levels among children is worrisome. As the world reopens and vaccines are distributed, it is important to determine whether these PA trends will persist and for how long. Following a natural disaster in East Japan in 2011, significantly decreased child PA levels persisted more than 3 years later.¹³² Consistently insufficient levels of PA in children around the world may have detrimental

effects on overall population health. The results of this review may be helpful in identifying barriers and facilitators to PA in children during the pandemic to determine what policies and programs would be most effective at increasing PA of children after the pandemic and beyond.

Summary – Accelerating Translation

The goal of "Effects of the COVID-19 Pandemic on Physical Activity in Children: A Systematic Review" was to synthesize the global impact of the COVID-19 pandemic on physical activity of children. Identifying changes in physical activity of children is important for understanding the extent to which the pandemic has affected already low levels of global physical activity among children. In addition, recognizing the biological, social, and environmental factors that affected changes in physical activity is critical for the development of effective programs to prevent short- and long-term health consequences. To the authors' knowledge, this is the first review to summarize the literature at the time of the data search regarding the global impact of the pandemic on the duration and types of physical activity children participated in and what biological, social, and environmental factors contributed to those changes. Preferred Reporting

Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines were used to conduct a database search on 03/29/2021. Included studies were peer-reviewed, in English language, and included both a measure of physical activity during the COVID-19 pandemic and physical activity data for children aged 18 and younger. The search yielded 673 unique citations and, ultimately, 69 articles were included in the review. The results of the study found that during the COVID-19 pandemic, frequently reported physical activity included walking, unstructured play, and virtual physical activity through online platforms. Of the articles that reported changes in physical activity of children during the COVID-19 pandemic, 89% reported physical activity (e.g., frequency, intensity, duration, or the percentage of children who met physical activity recommendations) of children decreased. Boys, younger children, children who lived with other children, and children with more outdoor play space had higher levels of physical activity. From these results, it was concluded that the COVID-19 pandemic resulted in decreased physical activity among children around the world. The results of this review may be helpful in identifying barriers and facilitators to physical activity in children during the COVID-19 pandemic. Future work in policy and program development is needed to target physical activity of children beyond the COVID-19 pandemic.

References

- Rundle AG, Park Y, Herbstman JB, Kinsey EW, Wang YC. COVID-19-related school closings and risk of weight gain among children. *Obesity* (Silver Spring). 2020;28(6):1008-9.
- World Health Organization. Coronavirus disease 2019 (COVID-19): situation report. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>. Last updated Jun 17, 2024; cited Jul 3, 2024.
- Nicola M, Alsafi Z, Sohrabi C, Kerwan A, Al-Jabir A, Iosifidis C, et al. The socio-economic implications of the coronavirus pandemic (COVID-19): a review. *Int J Surg*. 2020;78:185-93.
- Advisory Committee on Childhood Lead Poisoning Prevention, Centers for Disease Control and Prevention. Low level lead exposure harms children: a renewed call for primary prevention. Atlanta, GA: US Department of Health & Human Services, Centers for Disease Control and Prevention; 2012.
- Foster C, Moore JB, Singletary CR, Skelton JA. Physical activity and family-based obesity treatment: a review of expert recommendations on physical activity in youth. *Clin Obes*. 2018;8(1):68-79.
- Thivel D, Tremblay A, Genin PM, Panahi S, Riviere D, Duclos M. Physical activity, inactivity, and sedentary behaviors: definitions and implications in occupational health. *Front Public Health*. 2018;6:288.
- World Health Organization. Physical activity. Available from: <https://www.who.int/news-room/fact-sheets/detail/physical-activity>. Last updated Jun 26, 2024; cited Jul 3, 2024.
- Chaput JP, Willumsen J, Bull F, Chou R, Ekelund U, Firth J, et al. 2020 WHO guidelines on physical activity and sedentary behaviour for children and adolescents aged 5-17 years: summary of the evidence. *Int J Behav Nutr Phys Act*. 2020;17(1):141.
- Janssen I, Leblanc AG. Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *Int J Behav Nutr Phys Act*. 2010;7:40.
- Poitra VJ, Gray CE, Borghese MM, Carson V, Chaput JP, Janssen I, et al. Systematic review of the relationships between objectively measured physical activity and health indicators in school-aged children and youth. *Appl Physiol Nutr Metab*. 2016;41(6 Suppl 3):S197-239.
- Patterson R, McNamara E, Tainio M, de Sa TH, Smith AD, Sharp SJ, et al. Sedentary behaviour and risk of all-cause, cardiovascular and cancer mortality, and incident type 2 diabetes: a systematic review and dose response meta-analysis. *Eur J Epidemiol*. 2018;33(9):811-29.
- Guthold R, Stevens GA, Riley LM, Bull FC. Global trends in insufficient physical activity among adolescents: a pooled analysis of 298 population-based surveys with 1.6 million participants. *Lancet Child Adolesc Health*. 2020;4(1):23-35.
- Brazendale K, Beets MW, Weaver RG, Pate RR, Turner-McGrievy GM, Kaczynski AT, et al. Understanding differences between summer vs. school obesogenic behaviors of children: the structured days hypothesis. *Int J Behav Nutr Phys Act*. 2017;14(1):100.
- Franckle R, Adler R, Davison K. Accelerated weight gain among children during summer versus school year and related racial/ethnic disparities: a systematic review. *Prev Chronic Dis*. 2014;11:E101.
- Siebach MK, Piedimonte G, Ley SH. COVID-19 in childhood: Transmission, clinical presentation, complications and risk factors. *Pediatr Pulmonol*. 2021;56(6):1342-56.
- Dhochak N, Singhal T, Kabra SK, Lodha R. Pathophysiology of COVID-19: why children fare better than adults? *Indian J Pediatr*. 2020;87(7):537-46.
- Okuyama J, Seto S, Fukuda Y, Funakoshi S, Amai S, Onobe J, et al. Mental health and physical activity among children and adolescents during the COVID-19 pandemic. *Tohoku J Exp Med*. 2021;253(3):203-15.
- Ghosh R, Dubey MJ, Chatterjee S, Dubey S. Impact of COVID -19 on children: special focus on the psychosocial aspect. *Minerva Pediatr*. 2020;72(3):226-35.
- Shrestha R, Copenhaver M. Long-term effects of childhood risk factors on cardiovascular health during adulthood. *Clin Med Rev Vasc Health*. 2015;7:1-5.
- Kallio P, Pahkala K, Heinonen OJ, Tammelin TH, Palve K, Hirvensalo M, et al. Physical inactivity from youth to adulthood and adult cardiometabolic risk profile. *Prev Med*. 2021;145:106433.
- Rueggsegger GN, Booth FW. Health benefits of exercise. *Cold Spring Harb Perspect Med*. 2018;8(7):a029694.
- Lavie CJ, Ozemek C, Carbone S, Katzmarzyk PT, Blair SN. Sedentary behavior, exercise, and cardiovascular health. *Circ Res*. 2019;124(5):799-815.
- Booth FW, Roberts CK, Laye MJ. Lack of exercise is a major cause of chronic diseases. *Compr Physiol*. 2012;2(2):1143-211.
- Nieman DC, Wentz LM. The compelling link between physical activity and the body's defense system. *J Sport Health Sci*. 2019;8(3):201-17.
- Nieman DC, Pence BD. Exercise immunology: future directions. *J Sport Health Sci*. 2020;9(5):432-45.

26. Sallis R, Young DR, Tartof SY, Sallis JF, Sall J, Li Q, et al. Physical inactivity is associated with a higher risk for severe COVID-19 outcomes: a study in 48 440 adult patients. *Br J Sports Med.* 2021;55(19):1099-105.
27. Williams N, Radia T, Harman K, Agrawal P, Cook J, Gupta A. COVID-19 Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection in children and adolescents: a systematic review of critically unwell children and the association with underlying comorbidities. *Eur J Pediatr.* 2021;180(3):689-97.
28. Fernandes DM, Oliveira CR, Guerguis S, Eisenberg R, Choi J, Kim M, et al. Severe acute respiratory syndrome coronavirus 2 clinical syndromes and predictors of disease severity in hospitalized children and youth. *J Pediatr.* 2021;230:23-31 e10.
29. Gao YD, Ding M, Dong X, Zhang JJ, Kursat Azkur A, Azkur D, et al. Risk factors for severe and critically ill COVID-19 patients: a review. *Allergy.* 2021;76(2):428-55.
30. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ.* 2021;372:n71.
31. Yomoda K, Kurita S. Influence of social distancing during the COVID-19 pandemic on physical activity in children: a scoping review of the literature. *J Exerc Sci Fit.* 2021;19(3):195-203.
32. National Heart, Lung, Institute aB. Study quality assessment tools. Available from: <https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools>. Last updated Jul 2021; cited Jul 3, 2024.
33. Alonso-Martinez AM, Ramirez-Velez R, Garcia-Alonso Y, Izquierdo M, Garcia-Hermoso A. Physical activity, sedentary behavior, sleep and self-regulation in spanish preschoolers during the COVID-19 lockdown. *Int J Environ Res Public Health.* 2021;18(2):693.
34. Androutsos O, Perperidi M, Georgiou C, Chouliaras G. Lifestyle changes and determinants of children's and adolescents' body weight increase during the first COVID-19 lockdown in Greece: the COV-EAT study. *Nutrients.* 2021;13(3):930.
35. Cachon-Zagalaz J, Zagalaz-Sanchez Mf, Arufe-Giraldez V, Sanmiguel-Rodriguez A, Gonzalez-Valero G. Physical activity and daily routine among children aged 0-12 during the COVID-19 pandemic in Spain. *Int J Environ Res Public Health.* 2021;18(2):703.
36. Chambonniere C, Lambert C, Fearnbach N, Tardieu M, Fillon A, Genin P, et al. Effect of the COVID-19 lockdown on physical activity and sedentary behaviors in French children and adolescents: new results from the ONAPS national survey. *Eur J Integr Med.* 2021;43:101308.
37. Delisle Nystrom C, Alexandrou C, Henstrom M, Nilsson E, Okely AD, Wehbe El Masri S, et al. International study of movement behaviors in the early years (SUNRISE): results from SUNRISE Sweden's pilot and COVID-19 study. *Int J Environ Res Public Health.* 2020;17(22):8491.
38. Gilic B, Ostojic L, Corluka M, Volaric T, Sekulic D. Contextualizing parental/familial influence on physical activity in adolescents before and during COVID-19 pandemic: a prospective analysis. *Children (Basel).* 2020;7(9):125.
39. Gilic B, Zenic N, Separovic V, Jurcev Savicevic A, Sekulic D. Evidencing the influence of pre-pandemic sports participation and substance misuse on physical activity during the COVID-19 lockdown: a prospective analysis among older adolescents. *Int J Occup Med Environ Health.* 2021;34(2):151-63.
40. Sekulic D, Blazevic M, Gilic B, Kvesic I, Zenic N. Prospective analysis of levels and correlates of physical activity during COVID-19 pandemic and imposed rules of social distancing; gender specific study among adolescents from southern Croatia. *Sustainability.* 2020;12(10):4072.
41. Zenic N, Taiar R, Gilic B, Blazevic M, Maric D, Pojskic H, et al. Levels and changes of physical activity in adolescents during the COVID-19 pandemic: contextualizing urban vs. rural living environment. *Applied Sciences.* 2020;10(11):3997.
42. Francisco R, Pedro M, Delvecchio E, Espada JP, Morales A, Mazzeschi C, et al. Psychological symptoms and behavioral changes in children and adolescents during the early phase of COVID-19 quarantine in three European countries. *Front Psychiatry.* 2020;11:570164.
43. Hommes F, van Loon W, Thielecke M, Abramovich I, Lieber S, Hammerich R, et al. SARS-CoV-2 infection, risk perception, behaviour and preventive measures at schools in Berlin, Germany, during the early post-lockdown phase: a cross-sectional study. *Int J Environ Res Public Health.* 2021;18(5):2739.
44. Konstantinou C, Andrianou XD, Constantinou A, Perikkou A, Markidou E, Christophi CA, et al. Exposome changes in primary school children following the wide population non-pharmacological interventions implemented due to COVID-19 in Cyprus: a national survey. *EClinicalMedicine.* 2021;32:100721.
45. Kovacs VA, Starc G, Brandes M, Kaj M, Blagus R, Leskosek B, et al. Physical activity, screen time and the COVID-19 school closures in Europe - an observational study in 10 countries. *Eur J Sport Sci.* 2022;22(7):1094-103.
46. Lopez-Bueno R, Calatayud J, Andersen LL, Casana J, Ezzatvar Y, Casajus JA, et al. Cardiorespiratory fitness in adolescents before and after the COVID-19 confinement: a prospective cohort study. *Eur J Pediatr.* 2021;180(7):2287-93.
47. Lopez-Bueno R, Lopez-Sanchez GF, Casajus JA, Calatayud J, Gil-Salmeron A, Grabovac I, et al. Health-related behaviors among school-aged children and adolescents during the Spanish Covid-19 confinement. *Front Pediatr.* 2020;8:573.
48. Medrano M, Cadenas-Sanchez C, Osés M, Arenaza L, Amasene M, Labayen I. Changes in lifestyle behaviours during the COVID-19 confinement in Spanish children: a longitudinal analysis from the MUGI project. *Pediatr Obes.* 2021;16(4):e12731.
49. Lopez-Gil JF, Tremblay MS, Brazo-Sayavera J. Changes in healthy behaviors and meeting 24-h movement guidelines in Spanish and Brazilian preschoolers, children and adolescents during the COVID-19 lockdown. *Children (Basel).* 2021;8(2):83.
50. Morgul E, Kallitsoglou A, Essau CA. Psychological effects of the COVID-19 lockdown on children and families in the UK. *Revista de Psicologia Clinica con Ninos y Adolescentes.* 2020;7(3):42-8.
51. Poulain T, Meigen C, Sobek C, Ober P, Igel U, Korner A, et al. Loss of childcare and classroom teaching during the Covid-19-related lockdown in spring 2020: a longitudinal study on consequences on leisure behavior and schoolwork at home. *PLoS One.* 2021;16(3):e0247949.
52. Ng K, Cooper J, McHale F, Clifford J, Woods C. Barriers and facilitators to changes in adolescent physical activity during COVID-19. *BMJ Open Sport Exerc Med.* 2020;6(1):e000919.
53. Orgiles M, Morales A, Delvecchio E, Mazzeschi C, Espada JP. Immediate psychological effects of the COVID-19 quarantine in youth from Italy and Spain. *Front Psychol.* 2020;11:579038.
54. Pietrobelli A, Pecoraro L, Ferruzzi A, Heo M, Faith M, Zoller T, et al. Effects of COVID-19 lockdown on lifestyle behaviors in children with obesity living in Verona, Italy: a longitudinal study. *Obesity (Silver Spring).* 2020;28(8):1382-5.
55. Pombo A, Luz C, Rodrigues LP, Ferreira C, Cordovil R. Correlates of children's physical activity during the COVID-19 confinement in Portugal. *Public Health.* 2020;189:14-9.
56. Predieri B, Leo F, Candia F, Lucaccioni L, Madeo SF, Pugliese M, et al. Glycemic control improvement in Italian children and adolescents with type 1 diabetes followed through telemedicine during lockdown due to the COVID-19 pandemic. *Front Endocrinol (Lausanne).* 2020;11:595735.
57. Roe A, Blikstad-Balas M, Dalland CP. The impact of COVID-19 and homeschooling on students' engagement with physical activity. *Front Sports Act Living.* 2020;2:589227.
58. Ruiz-Roso MB, de Carvalho Padilha P, Matilla-Escalante DC, Brun P, Ulloa N, Acevedo-Correa D, et al. Changes of physical activity and ultra-processed food consumption in adolescents from different countries during Covid-19 pandemic: an observational study. *Nutrients.* 2020;12(8):2289.
59. Schmidt SCE, Anedda B, Burchartz A, Eichsteller A, Kolb S, Nigg C, et al. Physical activity and screen time of children and adolescents before and during the COVID-19 lockdown in Germany: a natural experiment. *Sci Rep.* 2020;10(1):21780.

60. Schmidt T, Pawlowski CS. Physical activity in crisis: the impact of COVID-19 on Danes' physical activity behavior. *Front Sports Act Living*. 2020;2:610255.
61. Salzano G, Passanisi S, Pira F, Sorrenti L, La Monica G, Pajno GB, et al. Quarantine due to the COVID-19 pandemic from the perspective of adolescents: the crucial role of technology. *Ital J Pediatr*. 2021;47(1):40.
62. Ten Velde G, Lubrecht J, Arayess L, van Loo C, Hesselink M, Reijnders D, et al. Physical activity behaviour and screen time in Dutch children during the COVID-19 pandemic: pre-, during- and post-school closures. *Pediatr Obes*. 2021;16(9):e12779.
63. Theis N, Campbell N, De Leeuw J, Owen M, Schenke KC. The effects of COVID-19 restrictions on physical activity and mental health of children and young adults with physical and/or intellectual disabilities. *Disabil Health J*. 2021;14(3):101064.
64. Tornaghi M, Lovecchio N, Vandoni M, Chirico A, Codella R. Physical activity levels across COVID-19 outbreak in youngsters of Northwestern Lombardy. *J Sports Med Phys Fitness*. 2021;61(7):971-6.
65. Wunsch K, Nigg C, Niessner C, Schmidt SCE, Oriwol D, Hanssen-Doose A, et al. The impact of COVID-19 on the interrelation of physical activity, screen time and health-related quality of life in children and adolescents in Germany: results of the Motorik-Modul study. *Children (Basel)*. 2021;8(2):98.
66. Zorcec T, Jakovska T, Micevska V, Boskovska K, Cholakovska VC. Pandemic with COVID-19 and families with children with chronic respiratory diseases. *Pril (Makedon Akad Nauk Umet Odd Med Nauki)*. 2020;41(2):95-101.
67. Carroll N, Sadowski A, Laila A, Hruska V, Nixon M, Ma DWL, et al. The impact of COVID-19 on health behavior, stress, financial and food security among middle to high income Canadian families with young children. *Nutrients*. 2020;12(8):2352.
68. Ellis WE, Dumas TM, Forbes LM. Physically isolated but socially connected: psychological adjustment and stress among adolescents during the initial COVID-19 crisis. *Can J Behav Sci*. 2020;52(3):177-87.
69. Guerrero MD, Vanderloo LM, Rhodes RE, Faulkner G, Moore SA, Tremblay MS. Canadian children's and youth's adherence to the 24-h movement guidelines during the COVID-19 pandemic: a decision tree analysis. *J Sport Health Sci*. 2020;9(4):313-21.
70. Hemphill NM, Kuan MTY, Harris KC. Reduced physical activity during COVID-19 pandemic in children with congenital heart disease. *Can J Cardiol*. 2020;36(7):1130-4.
71. McCormack GR, Doyle-Baker PK, Petersen JA, Ghoneim D. Parent anxiety and perceptions of their child's physical activity and sedentary behaviour during the COVID-19 pandemic in Canada. *Prev Med Rep*. 2020;20:101275.
72. Mitra R, Moore SA, Gillespie M, Faulkner G, Vanderloo LM, Chulak-Bozzer T, et al. Healthy movement behaviours in children and youth during the COVID-19 pandemic: exploring the role of the neighbourhood environment. *Health Place*. 2020;65:102418.
73. Moore SA, Faulkner G, Rhodes RE, Brussoni M, Chulak-Bozzer T, Ferguson LJ, et al. Impact of the COVID-19 virus outbreak on movement and play behaviours of Canadian children and youth: a national survey. *Int J Behav Nutr Phys Act*. 2020;17(1):85.
74. Bazett-Jones DM, Garcia MC, Taylor-Haas JA, Long JT, Rauh MJ, Paterno MV, et al. Impact of COVID-19 social distancing restrictions on training habits, injury, and care seeking behavior in youth long-distance runners. *Front Sports Act Living*. 2020;2:586141.
75. Dunton GF, Do B, Wang SD. Early effects of the COVID-19 pandemic on physical activity and sedentary behavior in children living in the U.S. *BMC Public Health*. 2020;20(1):1351.
76. Garcia JM, Lawrence S, Brazendale K, Leahy N, Fukuda D. Brief report: The impact of the COVID-19 pandemic on health behaviors in adolescents with autism spectrum disorder. *Disabil Health J*. 2021;14(2):101021.
77. McGuine TA, Biese KM, Petrovska L, Hetzel SJ, Reardon C, Kliethermes S, et al. Mental health, physical activity, and quality of life of US adolescent athletes during COVID-19-related school closures and sport cancellations: a study of 13 000 athletes. *J Athl Train*. 2021;56(1):11-9.
78. Pavlovic A, DeFina LF, Natale BL, Thiele SE, Walker TJ, Craig DW, et al. Keeping children healthy during and after COVID-19 pandemic: meeting youth physical activity needs. *BMC Public Health*. 2021;21(1):485.
79. Tulchin-Francis K, Stevens W, Jr., Gu X, Zhang T, Roberts H, Keller J, et al. The impact of the coronavirus disease 2019 pandemic on physical activity in U.S. children. *J Sport Health Sci*. 2021;10(3):323-32.
80. de Matos DG, Aida FJ, Almeida-Neto PFd, Moreira OC, Souza RFd, Marçal AC, et al. The impact of measures recommended by the government to limit the spread of coronavirus (COVID-19) on physical activity levels, quality of life, and mental health of Brazilians. *Sustainability*. 2020;12(21):9072.
81. Sa C, Pombo A, Luz C, Rodrigues LP, Cordovil R. Covid-19 social isolation in Brazil: effects on the physical activity routine of families with children. *Rev Paul Pediatr*. 2020;39:e2020159.
82. Siegle CBH, Pombo A, Luz C, Rodrigues LP, Cordovil R, Sa C. Influences of family and household characteristics on children's level of physical activity during social distancing due to Covid-19 in Brazil. *Rev Paul Pediatr*. 2020;39:e2020297.
83. Aguilar-Farias N, Toledo-Vargas M, Miranda-Marquez S, Cortinez-O'Ryan A, Cristi-Montero C, Rodriguez-Rodriguez F, et al. Sociodemographic predictors of changes in physical activity, screen time, and sleep among toddlers and preschoolers in Chile during the COVID-19 pandemic. *Int J Environ Res Public Health*. 2020;18(1):176.
84. Reece LJ, Owen K, Foley B, Rose C, Bellew B, Bauman A. Understanding the impact of COVID-19 on children's physical activity levels in NSW, Australia. *Health Promot J Austr*. 2021;32(2):365-6.
85. Cahal M, Amirav I, Diamant N, Be'er M, Besor O, Lavie M. Real-time effects of COVID-19 pandemic lockdown on pediatric respiratory patients. *Pediatr Pulmonol*. 2021;56(6):1401-8.
86. Chaturvedi K, Vishwakarma DK, Singh N. COVID-19 and its impact on education, social life and mental health of students: a survey. *Child Youth Serv Rev*. 2021;121:105866.
87. Ghanamah R, Eghbaria-Ghanamah H. Impact of COVID-19 pandemic on behavioral and emotional aspects and daily routines of Arab Israeli children. *Int J Environ Res Public Health*. 2021;18(6):2946.
88. Masi A, Mendoza Diaz A, Tully L, Azim SI, Woolfenden S, Efron D, et al. Impact of the COVID-19 pandemic on the well-being of children with neurodevelopmental disabilities and their parents. *J Paediatr Child Health*. 2021;57(5):631-6.
89. Munasinghe S, Sperandei S, Freebairn L, Conroy E, Jani H, Marjanovic S, et al. The impact of physical distancing policies during the COVID-19 pandemic on health and well-being among Australian adolescents. *J Adolesc Health*. 2020;67(5):653-61.
90. Nathan A, George P, Ng M, Wenden E, Bai P, Phiri Z, et al. Impact of COVID-19 restrictions on Western Australian children's physical activity and screen time. *Int J Environ Res Public Health*. 2021;18(5):2583.
91. Parker K, Uddin R, Ridgers ND, Brown H, Veitch J, Salmon J, et al. The use of digital platforms for adults' and adolescents' physical activity during the COVID-19 pandemic (Our Life at Home): survey study. *J Med Internet Res*. 2021;23(2):e23389.
92. Sciberras E, Patel P, Stokes MA, Coghill D, Middeldorp CM, Bellgrove MA, et al. Physical health, media use, and mental health in children and adolescents with ADHD during the COVID-19 pandemic in Australia. *J Atten Disord*. 2022;26(4):549-62.
93. Zhao Y, Guo Y, Xiao Y, Zhu R, Sun W, Huang W, et al. The effects of online homeschooling on children, parents, and teachers of grades 1-9 during the COVID-19 pandemic. *Med Sci Monit*. 2020;26:e925591.
94. World Health Organization. Adolescent health. Available from: <https://www.who.int/health-topics/adolescent-health>. Last updated Jun 10, 2024; cited Jul 3, 2024.
95. Wu X, Luo S, Zheng X, Ding Y, Wang S, Ling P, et al. Glycemic control in children and teenagers with type 1 diabetes around lockdown for COVID-19: a continuous glucose monitoring-based observational study. *J Diabetes Investig*. 2021;12(9):1708-17.
96. Zhang X, Zhu W, Kang S, Qiu L, Lu Z, Sun Y. Association between physical activity and mood states of children and adolescents in social isolation

- during the COVID-19 epidemic. *Int J Environ Res Public Health*. 2020;17(20):7666.
97. Abid R, Ammar A, Maaloul R, Souissi N, Hammouda O. Effect of COVID-19-related home confinement on sleep quality, screen time and physical activity in Tunisian boys and girls: a survey. *Int J Environ Res Public Health*. 2021;18(6):3065.
 98. Shinomiya Y, Yoshizaki A, Murata E, Fujisawa TX, Taniike M, Mohri I. Sleep and the general behavior of infants and parents during the closure of schools as a result of the COVID-19 pandemic: comparison with 2019 data. *Children (Basel)*. 2021;8(2):168.
 99. Xiang M, Zhang Z, Kuwahara K. Impact of COVID-19 pandemic on children and adolescents' lifestyle behavior larger than expected. *Prog Cardiovasc Dis*. 2020;63(4):531-2.
 100. Yang S, Guo B, Ao L, Yang C, Zhang L, Zhou J, et al. Obesity and activity patterns before and during COVID-19 lockdown among youths in China. *Clin Obes*. 2020;10(6):e12416.
 101. Elnaggar RK, Alqahtani BA, Mahmoud WS, Elfakharany MS. Physical activity in adolescents during the social distancing policies of the COVID-19 pandemic. *Asia Pac J Public Health*. 2020;32(8):491-4.
 102. Jia P, Zhang L, Yu W, Yu B, Liu M, Zhang D, et al. Impact of COVID-19 lockdown on activity patterns and weight status among youths in China: the COVID-19 Impact on Lifestyle Change Survey (COINLICS). *Int J Obes (Lond)*. 2021;45(3):695-9.
 103. Prince SA, Butler GP, Rao DP, Thompson W. Evidence synthesis - where are children and adults physically active and sedentary? - A rapid review of location-based studies. *Health Promot Chronic Dis Prev Can*. 2019;39(3):67-103.
 104. Kelso A, Reimers AK, Abu-Omar K, Wunsch K, Niessner C, Wasche H, et al. Locations of physical activity: where are children, adolescents, and adults physically active? A systematic review. *Int J Environ Res Public Health*. 2021;18(3):1240.
 105. Weaver RG, Beets MW, Perry M, Hunt E, Brazendale K, Decker L, et al. Changes in children's sleep and physical activity during a 1-week versus a 3-week break from school: a natural experiment. *Sleep*. 2019;42(1):zsy205.
 106. Marzi I, Demetriou Y, Reimers AK. Social and physical environmental correlates of independent mobility in children: a systematic review taking sex/gender differences into account. *Int J Health Geogr*. 2018;17(1):24.
 107. Steene-Johannessen J, Hansen BH, Dalene KE, Kolle E, Northstone K, Moller NC, et al. Variations in accelerometry measured physical activity and sedentary time across Europe - harmonized analyses of 47,497 children and adolescents. *Int J Behav Nutr Phys Act*. 2020;17(1):38.
 108. D'Angelo H, Fowler SL, Nebeling LC, Oh AY. Adolescent physical activity: moderation of individual factors by neighborhood environment. *Am J Prev Med*. 2017;52(6):888-94.
 109. Alharbi M. Influence of individual and family factors on physical activity among Saudi girls: a cross-sectional study. *Ann Saudi Med*. 2019;39(1):13-21.
 110. Lu C, Stolk RP, Sauer PJ, Sijtsma A, Wiersma R, Huang G, et al. Factors of physical activity among Chinese children and adolescents: a systematic review. *Int J Behav Nutr Phys Act*. 2017;14(1):36.
 111. Reimers AK, Schmidt SCE, Demetriou Y, Marzi I, Woll A. Parental and peer support and modelling in relation to domain-specific physical activity participation in boys and girls from Germany. *PLoS One*. 2019;14(10):e0223928.
 112. Best K, Ball K, Zarnowiecki D, Stanley R, Dollman J. In search of consistent predictors of children's physical activity. *Int J Environ Res Public Health*. 2017;14(10):1258.
 113. Dishman RK, McIver KL, Dowda M, Saunders RP, Pate RR. Self-efficacy, beliefs, and goals: moderation of declining physical activity during adolescence. *Health Psychol*. 2019;38(6):483-93.
 114. Burns RD, Bai Y, Fu Y, Pfledderer CD, Brusseau TA. Parent engagement and support, physical activity, and academic performance (PESPAAP): a proposed theoretical model. *Int J Environ Res Public Health*. 2019;16(23):4698.
 115. Khan SR, Uddin R, Mandic S, Khan A. Parental and peer support are associated with physical activity in adolescents: evidence from 74 countries. *Int J Environ Res Public Health*. 2020;17(12):4435.
 116. Haidar A, Ranjit N, Archer N, Hoelscher DM. Parental and peer social support is associated with healthier physical activity behaviors in adolescents: a cross-sectional analysis of Texas School Physical Activity and Nutrition (TX SPAN) data. *BMC Public Health*. 2019;19(1):640.
 117. Cooper AR, Goodman A, Page AS, Sherar LB, Esliger DW, van Sluijs EM, et al. Objectively measured physical activity and sedentary time in youth: the International Children's Accelerometry Database (ICAD). *Int J Behav Nutr Phys Act*. 2015;12:113.
 118. Craggs C, Corder K, van Sluijs EM, Griffin SJ. Determinants of change in physical activity in children and adolescents: a systematic review. *Am J Prev Med*. 2011;40(6):645-58.
 119. Hu D, Zhou S, Crowley-McHattan ZJ, Liu Z. Factors that influence participation in physical activity in school-aged children and adolescents: a systematic review from the social ecological model perspective. *Int J Environ Res Public Health*. 2021;18(6):3147.
 120. Corder K, Sharp SJ, Atkin AJ, Griffin SJ, Jones AP, Ekelund U, et al. Change in objectively measured physical activity during the transition to adolescence. *Br J Sports Med*. 2015;49(11):730-6.
 121. Pate RR, Sallis JF, Ward DS, Stevens J, Dowda M, Welk GJ, et al. Age-related changes in types and contexts of physical activity in middle school girls. *Am J Prev Med*. 2010;39(5):433-9.
 122. Rhodes RE, Guerrero MD, Vanderloo LM, Barbeau K, Birken CS, Chaput JP, et al. Development of a consensus statement on the role of the family in the physical activity, sedentary, and sleep behaviours of children and youth. *Int J Behav Nutr Phys Act*. 2020;17(1):74.
 123. Edwards MJ, Jago R, Sebire SJ, Kesten JM, Pool L, Thompson JL. The influence of friends and siblings on the physical activity and screen viewing behaviours of children aged 5-6 years: a qualitative analysis of parent interviews. *BMJ Open*. 2015;5(5):e006593.
 124. Loebach J, Sanches M, Jaffe J, Elton-Marshall T. Paving the way for outdoor play: examining socio-environmental barriers to community-based outdoor play. *Int J Environ Res Public Health*. 2021;18(7):3617.
 125. Euler R, Jimenez EY, Sanders S, Kuhlemeier A, Van Horn ML, Cohen D, et al. Rural-urban differences in baseline dietary intake and physical activity levels of adolescents. *Prev Chronic Dis*. 2019;16:E01.
 126. Rivera-Ochoa M, Brazo-Sayavera J, Vizmanos-Lamotte B, Manas A, Lopez-Taylor JR, Gonzalez-Gross M, et al. Health-related factors in rural and urban Mexican adolescents from the State of Jalisco: the HELENA-MEX study. *Int J Environ Res Public Health*. 2020;17(23):8959.
 127. Adebusoje B, Phalkey R, Leonardi-Bee J, Chattopadhyay K. Association of the built environment with physical activity in children and adolescents in Africa: a systematic review protocol. *JBI Evid Synth*. 2020;18(3):553-63.
 128. Gray C, Gibbons R, Larouche R, Sandseter EB, Bienenstock A, Brussoni M, et al. What is the relationship between outdoor time and physical activity, sedentary behaviour, and physical fitness in children? A systematic review. *Int J Environ Res Public Health*. 2015;12(6):6455-74.
 129. Larouche R, Garriguet D, Gunnell KE, Goldfield GS, Tremblay MS. Outdoor time, physical activity, sedentary time, and health indicators at ages 7 to 14: 2012/2013 Canadian Health Measures Survey. *Health Rep*. 2016;27(9):3-13.
 130. Lindsay AC, Greaney ML, Wallington SF, Mesa T, Salas CF. A review of early influences on physical activity and sedentary behaviors of preschool-age children in high-income countries. *J Spec Pediatr Nurs*. 2017;22(3):e12182.
 131. Christiana RW, Bouldin ED, Battista RA. Active living environments mediate rural and non-rural differences in physical activity, active transportation, and screen time among adolescents. *Prev Med Rep*. 2021;23:101422.
 132. Okazaki K, Suzuki K, Sakamoto Y, Sasaki K. Physical activity and sedentary behavior among children and adolescents living in an area affected by the 2011 Great East Japan earthquake and tsunami for 3 years. *Prev Med Rep*. 2015;2:720-4.

Acknowledgments

The authors acknowledge Brianna Andre and Steven Moore from the Sladen Library at Henry Ford Hospital for their assistance with the literature search. The authors acknowledge Abigail C. Radomsky, a 4th year medical student from Wayne State University School of Medicine, for her assistance with abstract review and article selection.

Conflict of Interest Statement & Funding

The Authors have no funding, financial relationships or conflicts of interest to disclose.

Author Contributions

Conceptualization, A.R.; Investigation, A.R., A.C.R.; Writing – Original Draft, A.R.; Writing – Review and Editing, A.B.R., S.S.; Visualization, A.R.; Supervision, A.B.R., S.S.

Cite as

Ramirez A, Rapp AB, Santarossa S. Effects of the COVID-19 Pandemic on Physical Activity in Children: A Systematic Review. Int J Med Stud. 2024 Jul-Sep;12(3):311-333.

This work is licensed under a [Creative Commons Attribution 4.0 International License](#)

ISSN 2076-6327

This journal is published by [Pitt Open Library Publishing](#)



Supplementary Material

Literature search results for Systematic Review on Covid-19 and Physical Activity in Children, 2020-present, English

Searches ran on March 29, 2021

Ovid MEDLINE: 402

Ovid MEDLINE(R) and Epub Ahead of Print, In-Process, In-Data-Review & Other Non-Indexed Citations, Daily and Versions(R) <1946 to March 26, 2021>

1 (exp Exercise/ or exp Sedentary Behavior/ or Dancing/ or exp "play and playthings"/ or exp Sports/ or Screen Time/ or (exercise or exercises or sedentary or screen-time or screen-use or fitness or (physical adj2 (activit* or inactivity or education))).ti,ab,kf.) and (exp adolescent/ or exp child/ or (adolescen* or boy? or boyfriend or boyhood or girlfriend or girlhood or child* or girl? or juvenil* or kid? or minors or minors* or paediatric* or paediatric* or pediatric* or preschool* or puber* or pubescen* or school* or teen* or underage? or under-age? or youth*).ti,ab,kf. or (pediatric* or paediatric* or child* or adolescen* or young).jn,jw.) 135600
2 limit 1 to (english language and yr="2020 -Current" and covid-19) 402

Embase Results: 263

((('exercise'/exp/mj OR 'sedentary lifestyle'/de OR 'physical activity'/exp OR 'play'/exp OR 'sport'/exp/mj OR 'screen time'/de OR 'physical inactivity'/exp OR 'sitting'/de OR 'sedentary time'/de OR 'youth sport'/de) OR ('fitness':ti,ab OR 'sedentary':ti,ab OR 'screen time':ti,ab OR 'screen use':ti,ab) OR (physical NEAR/2 (activit* OR inactivity)):ti,ab) AND (('juvenile'/mj OR 'boy'/mj OR 'girl'/mj OR 'child'/mj OR 'preschool child'/mj OR 'school child'/mj OR 'adolescent'/exp/mj OR 'minor (person)'/exp/mj OR 'adolescence'/exp/mj OR 'childhood'/exp/mj OR boy:ti,ab OR boy:ti,ab OR boyfriend:ti,ab OR boyhood:ti,ab OR girl*:ti,ab OR kid:ti,ab OR kids:ti,ab OR child:ti,ab OR child*:ti,ab OR children*:ti,ab OR schoolchild*:ti,ab OR schoolchild:ti,ab OR 'school child':ti,ab OR 'school child*':ti,ab OR adolescen*:ti,ab OR juvenil*:ti,ab OR youth*:ti,ab OR teen*:ti,ab OR under*age*:ti,ab OR pubescen*:ti,ab) AND ('coronavirus disease 2019'/exp OR 'severe acute respiratory syndrome coronavirus 2'/exp OR 'severe acute respiratory syndrome coronavirus 2' OR '2019 ncov' OR 2019ncov OR 'covid-19' OR '2019-ncov' OR 'sars-cov-2' OR '2019ncov' OR 'novel coronavirus') AND [english]/lim AND [2020-2021]/py 263

PsycInfo: 39

APA PsycInfo <2002 to March Week 4 2021>

(exp "Summer Camps (Recreation)"/ or exp Leisure Time/ or exp Exercise/ or exp Physical Activity/ or Activity Level/ or Physical Fitness/ or exp Sports/ or Athletic Participation/ or Dance/ or Childhood Play Behavior/ or Childrens Recreational Games/ or (exercise or exercises or sedentary or screen-time or screen-use or fitness or (physical adj2 (activit* or inactivity))).ti,ab.) and (minors or minors* or boy or boys or boyfriend or boyhood or girl* or kid or kids or child or child* or children* or schoolchild* or schoolchild or school child or school child* or adolescen* or juvenil* or youth* or teen* or under*age* or pubescen* or pediatric* or paediatric* or paediatric* or school or school*).mp. and (coronavirus/ or ('severe acute respiratory syndrome coronavirus 2' or '2019 ncov' or 2019ncov or 'covid-19' or '2019-ncov' or 'sars-cov-2' or '2019ncov' or 'novel coronavirus').mp.) 62
limit 9 to (english language and yr="2020 -Current") 39

CINAHL: 168

S9 S4 AND S5 AND S6 Limiters - Published Date: 20200101-20211231
Narrow by Language: - English 168
S8 S4 AND S5 AND S6 Limiters - Published Date: 20200101-20211231 168
S7 S4 AND S5 AND S6 179
S6 (MH "COVID-19") OR (MH "Coronavirus Infections+") OR (MH "Coronavirus+") OR (TX 'severe acute respiratory syndrome coronavirus 2' or '2019 ncov' or 2019ncov or 'covid-19' or '2019-ncov' or 'sars-cov-2' or '2019ncov' or 'novel coronavirus') 48,402
S5 (MH "Hospitals, Pediatric") OR (MH "Adolescence+") OR (MH "Maternal Age 14 and Under") OR (MH "Child, Preschool") OR (MH "Child Psychology") OR (MH "Pediatric Units") OR (MH "Intensive Care Units, Pediatric") OR (MH "Rehabilitation, Pediatric") OR (TX minors or minors* or boy or boys or boyfriend or boyhood or girl* or kid or kids or child or child* or children* or schoolchild* or schoolchild or school child or school child* or adolescen* or juvenil* or youth* or teen* or under*age* or pubescen* or pediatric* or paediatric* or paediatric*) 1,237,921
S4 S1 OR S2 OR S3 380,492
S3 TX exercise or exercises or sedentary or "screen time" or "screen use" or fitness or "physical activity" or "physical inactivity" 302,701
S2 (MH "Life Style, Sedentary+") 8,845
S1 (MH "Physical Activity") OR (MH "Physical Fitness+") OR (MH "Exercise+") OR (MH "Dancing+") OR (MH "Play and Playthings+") OR (MH "Sports+") OR (MH "Life Style, Sedentary") OR (MH "Screen Time") 242,545

Web of Science: 168

TS=(("severe acute respiratory syndrome coronavirus 2" or "2019 ncov" or 2019ncov or "covid-19" or "2019-ncov" or "sars-cov-2" or "2019ncov" or "novel coronavirus")

AND

TS=(minors or minors* or boy or boys or boyfriend or boyhood or girl* or kid or kids or child or child* or children* or schoolchild* or schoolchild or school child or school child* or adolescen* or juvenil* or youth* or teen* or under*age* or pubescen* or pediatric* or paediatric* or paediatric*)

AND

TS=(exercise OR sedentary OR "physical activity" OR "physical inactivity" OR "activity level" OR sports OR "screen time" OR "screen use" OR fitness)

Indexes=SCI-EXPANDED Timespan=All years

Total results = 1040

Total duplicates = 367

Total after deduplicating = 673