

1 **Title:** A Case Report of Bardet Biedl Syndrome in a Patient from Pakistan who Presented with Osmotic  
2 Symptoms associated with Diabetes Mellitus.

3 **Article type:** Case Report

4 **Author names:**

- 5 1. Bakht Babar
- 6 2. Muqsit Ali Shaukat
- 7 3. Misbah Manzoor
- 8 4. Samina Bibi
- 9 5. Hashim Khan

10 **Degrees and Affiliations:**

- 11 1. FCPS Medicine, Fellow and Registrar Endocrinology. Hayatabad Medical Complex, Peshawar, Pakistan.
- 12 2. MBBS. Resident Physician in General Medicine at Federal Government Polyclinic Hospital Islamabad,  
13 Pakistan.
- 14 3. FCPS Cardiology, Registrar. Peshawar Institute of Cardiology, Peshawar, Pakistan.
- 15 4. FCPS Medicine, Fellow Endocrinology. Hayatabad Medical Complex, Peshawar, Pakistan.
- 16 5. Final-year Medical student. Peshawar Medical College, Peshawar, Pakistan.

17  
18 **ORCID (Open Researcher and Contributor Identifier):**

- 19 1. <https://orcid.org/0000-0002-5491-2393>
- 20 2. <https://orcid.org/0000-0002-9961-3354>
- 21 3. <https://orcid.org/0000-0001-5615-8382>
- 22 4. <https://orcid.org/0000-0001-8382-861X>
- 23 5. <https://orcid.org/0000-0003-2625-3201>

24 **Corresponding author email:** muqsitali@hotmail.com

25 **Acknowledgment:** None

26 **Financing:** No grant, financial support or any other support was made to the study or the authors from any  
27 source.

28 **Conflict of interest statement by authors:** There is no conflict of interest for this publication.

29 **Compliance with ethical standards:** Informed consent was obtained from the patient's representative.

30 **Authors Contribution Statement:** Conceptualization: B.B., M.A.S., M.M., S.B.. Investigation: H.K.. Project  
31 Administration: B.B.. Writing - Original Draft: B.B., M.A.S., M.M.. Writing - Review Editing: B.B., M.A.S..

32 **Highlights**

- 33 ● Patients presenting with polyuria, polydipsia, polyphagia along with the presence of syndromic features,  
34 should be assessed for BBS.
- 35 ● The treatment options available treat diabetes mellitus in BBS include weight loss, exercise, oral  
36 hypoglycemic drugs, and insulin

37 **Manuscript word count:** 1359

38 **Abstract word count:** 238

39 **Number of Figures and Tables:** 2 Figures and 1 table.

40

1 **Discussion Points:** Bardet Biedl Syndrome (BBS) is rare disorder inherited in an autosomal recessive pattern.  
2 This article describes a case of BBS and emphasis upon the management of diabetes in BBS.

3  
4

5 **Dates**

6 Submission: 12/12/2022

7 Revisions: 04/01/2023, 01/10/2024

8 Responses: 05/17/2023, 03/17/2024

9 Acceptance: 06/20/2024

10 Publication: 06/27/2024

11

12 **Editors**

13 Associate Editor/Editor: Francisco J. Bonilla-Escobar

14 Student Editors: Alisha Poppen, L V Simhachalam Kutikuppala

15 Copyeditor:

16 Proofreader:

17 Layout Editor:

18

19 **Publisher's Disclosure:** *This is a PDF file of an unedited manuscript that has been accepted for publication.*  
20 *As a service to our readers and authors we are providing this early version of the manuscript. The manuscript*  
21 *will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable*  
22 *form. Please note that during the production process errors may be discovered which could affect the content,*  
23 *and all legal disclaimers that apply to the journal pertain.*

24

25

1 **ABSTRACT.**

2

3 **Background:** The primary features of Bardet Biedl syndrome (BBS) are characterized by retinal  
4 degeneration, central obesity, post-axial polydactyly, intellectual impairment, hypogonadism, and renal  
5 anomalies. The secondary clinical features are syndactyly, delay in the achievement of developmental  
6 milestones, diabetes insipidus, diabetes mellitus, congenital heart disease and liver involvement. It is  
7 diagnosed when a patient has either four primary or three primary and two secondary clinical features and the  
8 management depends upon those predominant features.

9

10 **Case presentation:** A 12-year-old boy presented with polyuria, polydipsia and polyphagia. His developmental  
11 milestones were delayed and had syndromic features on physical examination. Ultrasound of the abdomen and  
12 pelvis showed mild right-side hydronephrosis. His BMI was 28 and HbA1c was 18.3%, and values of LH and  
13 FSH were decreased. As a result, he was diagnosed with a case of BBS. He was commenced on insulin  
14 therapy with Insulin Regular (Human Insulin) and Insulin NPH (Human Insulin).

15

16 **Conclusion:** Patients presenting with polyuria, polydipsia, polyphagia along with the presence of syndromic  
17 features, should be assessed for BBS as osmotic symptoms associated with diabetes mellitus can be initial  
18 clinical presentation of BBS. Its management depends upon the signs and symptoms of the patient. Various  
19 options including weight loss, exercise, oral hypoglycemic drugs, and insulin are available to treat diabetes  
20 mellitus in BBS. Some cases can have very high levels of HbA1c on initial presentation, which may require  
21 insulin without trying oral hypoglycemic drugs to treat diabetes mellitus in BBS.

22

23 **Key Words:** bardet biedl syndrome, bardet-biedl syndrome, laurence moon bardet biedl syndrome, diabetes  
24 mellitus,

25

1 **INTRODUCTION.**

2

3 Bardet Biedl syndrome (BBS) is an autosomal recessive ciliopathy, with a prevalence of 1 in 13,500 to  
4 160,000 people.<sup>1</sup> It has primary as well as some secondary clinical features. The primary clinical features of  
5 BBS include retinal degeneration in 94%, central obesity in 89%, post-axial polydactyly in 79%, intellectual  
6 impairment in 66%, hypogonadism in 59%, and renal anomalies in 52% of patients.<sup>7</sup> Secondary clinical  
7 features of BBS include syndactyly, delay in the achievement of developmental milestones, diabetes insipidus,  
8 diabetes mellitus, and congenital heart disease.<sup>3</sup> It is diagnosed when a patient has either four primary or  
9 three primary and two secondary clinical features.<sup>7</sup> It is managed according to the signs and symptoms of the  
10 patient and requires a multidisciplinary approach.<sup>9</sup>

11

12 We represent a case of BBS in a 12-year-old male, newly diagnosed diabetic patient who presented with  
13 polyuria, polydipsia and polyphagia. He had all the six primary clinical features of BBS. Among secondary  
14 clinical features of BBS he had syndactyly, developmental delay and diabetes mellitus. The presence of  
15 osmotic symptoms of diabetes mellitus as a predominant feature of BBS and management of diabetes  
16 mellitus in BBS has been rarely discussed in the literature.<sup>5</sup>

17

18

Accepted, in

**1 THE CASE**

2

3 A 12-year-old boy presented to the outpatient department of Hayatabad Medical Complex, Peshawar,  
4 Pakistan complaining of polyuria, polydipsia, and polyphagia. According to the patient's mother, he was  
5 obese, had polyphagia and a low intelligence quotient (IQ) since childhood. For the past one and a half  
6 months, he had increased frequency of urination, excessive thirst and increased appetite. It was further  
7 revealed that he was born out of a non-consanguineous marriage, through a normal vaginal delivery and was  
8 fully immunized. He started walking at 3 years of age and started speaking at 4 years of age. He was dropped  
9 out of school because of learning difficulties at the school. On examination, he had syndromic features  
10 including bilateral post-axial polydactyly in both of his feet (Figure 1), syndactyly in his right hand (Figure 2),  
11 widely spaced eyes, depressed nasal bridge, and high-pitched sound. His body mass index (BMI) was 28.  
12 Head, ears, eyes, nose, and throat examination revealed bilateral conductive deafness and features of retinitis  
13 pigmentosa on fundoscopy. Central nervous system examination revealed no sensory and motor  
14 abnormalities. Genital examination revealed a small penis (<1cm), small and soft testis, and a redundant  
15 scrotum that was difficult to accommodate. The rest of the examination was unremarkable. Laboratory  
16 investigations done in the hospital are given below in Table 1.

17

18 Urinalysis and urine microscopy showed no abnormal laboratory values. An echocardiography was performed  
19 which was normal with no structural defects. Ultrasound of the abdomen and pelvis showed mildly enlarged  
20 spleen and mild right-side hydronephrosis. He was started on an injection of Insulin Regular (Human Insulin) 2  
21 units three times a day and an injection of Insulin NPH (Human Insulin) 2 units in the morning and 4 units at  
22 night for the management of his high blood sugar.

23

Accepted

1 **DISCUSSION.**

2

3 The prevalence of BBS ranges from 1 in 13,500 in some Arabian countries like Kuwait to 1 in 1,60,000 in  
4 northern Europe. In Pakistan, the exact data regarding the prevalence of BBS is not available. Very few case  
5 reports have been reported of this disorder in Pakistan.<sup>2</sup> We report a case of BBS in a 12-year-old male  
6 patient from Peshawar, Pakistan who presented with polyuria, polydipsia, and polyphagia. He was diagnosed  
7 with diabetes mellitus and was later found to have met the clinical criteria for BBS.

8

9 The diagnosis of BBS is made on the basis of clinical signs and symptoms. It has six primary or cardinal  
10 features as well as some secondary features. The primary features of this syndrome include cone-rod  
11 dystrophy, obesity, hypogonadism, intellectual impairment, renal abnormalities, and polydactyl.<sup>3,4</sup> Our patient  
12 had all of these clinical features. The presence of all the primary features of BBS in a single patient is rare.<sup>5</sup>  
13 Cone-rod dystrophy manifested in the form of retinitis pigmentosa on retinal screening, he was overweight as  
14 his BMI was 28, hypogonadism appeared in the form of small penis, small and soft testis, redundant scrotum,  
15 low levels of LH and FSH and low to normal level of serum testosterone, he was dropped out of school due to  
16 learning difficulties, renal involvement occurred in the form of right sided hydronephrosis and he had an extra  
17 digit in his both feet. Secondary features of BBS include, syndactyly, delay in the achievement of  
18 developmental milestones, diabetes insipidus, diabetes mellitus, and congenital heart disease.<sup>3</sup> Our patient  
19 had syndactyly in his right hand. He had developmental delay as he started walking at the age of 3 and  
20 speaking at age 4. He was newly diagnosed with diabetes as his random blood sugar was 360 mg/dl and  
21 HbA1c was 18.3%.

22

23 Our patient had dysmorphic facial features including widely spaced eyes, a depressed nasal bridge, and a  
24 long philtrum. These features have been described to be present in cases of BBS. Other facial features of  
25 BBS described in the literature but not present in our patient include high arched palate, less number of teeth,  
26 and retrognathia.<sup>4,6</sup>

27

28 BBS is inherited in an autosomal recessive form. Mutations twenty one different subtypes in the BBS gene  
29 can be responsible for BBS. Since BBS is diagnosed clinically by the presence of either four primary clinical  
30 features or three primary and two secondary clinical features and it doesn't require the confirmation of the  
31 presence of genetic mutation, we did not do genetic testing in our patient.<sup>7</sup> Moreover, genetic testing was not  
32 feasible as well in our resource limited setting.

33

34 Very few cases of BBS with diabetes mellitus have been reported in the literature to the best of our  
35 knowledge. Quader et al.<sup>8</sup> described a case of Laurence-Moon-Bardet-Biedl syndrome (LMBBS) in a diabetic  
36 patient who presented with loss of vision, obesity, developmental delay and learning difficulties. Our patient  
37 was different than that as he initially presented with the osmotic symptoms of diabetes mellitus like polyuria,  
38 polydipsia, and polyphagia. On further evaluation, he was later on found to have primary and secondary  
39 features of BBS. This initial clinical presentation of BBS with osmotic symptoms is described in very few cases  
40 in the literature. Garg et al. also described a case of LMBBS with diabetes in a 22-year-old male patient.

1 Though he presented initially with osmotic symptoms of polyuria and polydipsia, it was described to be an  
2 incomplete manifestation of LMBBS due to the lack of digital anomalies.<sup>5</sup>

3  
4 The management of BBS requires a multidisciplinary approach, keeping in view the predominant signs and  
5 symptoms of the patient.<sup>9</sup> The predominant features in our patient were osmotic symptoms of diabetes  
6 mellitus. Diabetes mellitus in patients with BBS can be diagnosed using the guidelines of the American  
7 Diabetes Association (ADA).<sup>10</sup> A case-control study conducted by Mujahid et al. has shown that the patients  
8 with BBS are two times more likely to develop insulin resistance.<sup>11</sup> Therefore, the effective ways to treat  
9 diabetes mellitus in BBS could include exercise, weight loss, and insulin sensitizers including metformin and  
10 thiazolidinediones.<sup>12</sup> But few cases of BBS have been reported which showed resistance to oral  
11 hypoglycemic drugs as well and were ultimately started on insulin. A case report of BBS with diabetes mellitus  
12 was reported by garg et al. in which the patient was resistant to oral hypoglycemic drugs and was  
13 consequently started on insulin.<sup>5</sup> Quader et al. also started a 13-year-old boy having BBS on metformin for the  
14 control of his diabetes mellitus but the data regarding his follow up and how effectively it controlled his  
15 diabetes mellitus was not given.<sup>8</sup> Our patient had very high level of HbA1c. The type of diabetes mellitus  
16 which affects the patients with BBS is mostly reported to be type 2 diabetes mellitus<sup>4</sup> and insulin  
17 administration is recommended for people whose HbA1C is greater than 10% or blood glucose levels more  
18 than 300mg/dl. <sup>13</sup> Apart from that, few patients have been reported to have no response to oral hypoglycemic  
19 drugs.<sup>5</sup> Due to these reasons, we started the patient on a basal bolus regimen.

20 The prognosis of BBS is affected by multiple factors. Patients may suffer functional impairment from  
21 progressive decrease in vision. In BBS, the leading cause of death is kidney disease but the life expectancy  
22 has been reported to be normal.<sup>14</sup> We arranged a follow up meeting with our patient to see the response to  
23 treatment and monitor the long term effects of BBS but he did not show up for follow up.

24

Accepted

1 **REFERENCES.**

- 2
- 3 1. Forsythe E, Beales PL. Bardet–Biedl syndrome. *Eur J Hum Genet.* 2013;21(1):8–13.
- 4 2. Khan OA, Majeed R, Saad M, Khan A, Ghassan A. Rarity of Laurence Moon Bardet Biedl  
5 Syndrome and its Poor Management in the Pakistani Population. *Cureus.* 2019;11(2):e4114.
- 6 3. Medscape. A Retinal Disease With Systemic Associations. Available from:  
7 <https://www.medscape.com/viewarticle/479108>. Last updated June 10, 2004; cited Dec 4, 2022 .
- 8 4. NORD (National Organization for Rare Disorders). Bardet-Biedl Syndrome. Available from:  
9 <https://rarediseases.org/rare-diseases/bardet-biedl-syndrome/>. Last Updated July 12, 2022; cited  
10 Dec 4, 2022.
- 11 5. GARG M, MADHU S, DWARAKANATH C, AMMINI A. LAURENCE-MOON-BARDET-BIEDL  
12 SYNDROME – PRESENTING WITH ACUTE ONSET OF DIABETES MELLITUS. *Med J Armed  
13 Forces India.* 1998;54(2):155–6.
- 14 6. Forsythe E, Kenny J, Bacchelli C, Beales PL. Managing Bardet-Biedl Syndrome-Now and in the  
15 Future. *Front Pediatr.* 2018;6:23.
- 16 7. National Institutes of Health. Bardet-Biedl Syndrome Overview. Available from:  
17 <http://www.ncbi.nlm.nih.gov/books/NBK1363/>. Last Updated March 23, 2023: cited June 14, 2024.
- 18 8. Quader M, Islam MS, Ahmed QMU, Azad MAK, Rahim MA. Laurence -Moon-Bardet- Biedl  
19 syndrome with Diabetes Mellitus. *Journal of Medicine.* 2012;13(1):97–9.
- 20 9. Caba L, Florea L, Braha EE, Lupu VV, Gorduza EV. Monitoring and Management of Bardet-Biedl  
21 Syndrome: What the Multi-Disciplinary Team Can Do. *JMDH.* 2022;15:2153–67.
- 22 10. ADA. Diabetes Diagnosis and Tests. Available from: <https://diabetes.org/diabetes/a1c/diagnosis>.  
23 Dec 4, cited 2022.
- 24 11. Mujahid S, Hunt KF, Cheah YS, Forsythe E, Hazlehurst JM, Sparks K, et al. The Endocrine and  
25 Metabolic Characteristics of a Large Bardet-Biedl Syndrome Clinic Population. *J Clin Endocrinol  
26 Metab.* 2018;103(5):1834–41.
- 27 12. ADA. Insulin Resistance. Available from: [https://diabetes.org/healthy-living/medication-  
28 treatments/insulin-resistance](https://diabetes.org/healthy-living/medication-treatments/insulin-resistance). cited Dec 4, 2022.
- 29 13. ElSayed NA, Aleppo G, Aroda VR, Bannuru RR, Brown FM, Bruemmer D, et al. 9. Pharmacologic  
30 Approaches to Glycemic Treatment: Standards of Care in Diabetes—2023. *Diabetes Care.*  
31 2022;46(Supplement\_1):S140–57.
- 32 14. Myriad Women’s Health. Bardet-Biedl syndrome, BBS12-related. Available from:  
33 <https://myriad.com/womens-health/diseases/bardet-biedl-syndrome-bbs12-related/>. cited May 16,  
34 2023.
- 35
- 36
- 37



**1 SUMMARY - ACCELERATING TRANSLATION**

2 Bardet Biedl syndrome (BBS) is a rare disorder which can affect eyes, kidneys, and can cause weight gain  
3 and intellectual impairment. It can also cause digital anomalies resulting in extra digits or fused digits. Apart  
4 from that developmental milestones can be delayed and a patient can have diabetes mellitus as well. This  
5 article describes a case of BBS in a 12-years-old boy and emphasizes the management of high blood sugar  
6 levels in BBS. He presented with increased frequency of urination, increased thirst, and increased appetite.  
7 His developmental milestones were delayed, he has low intelligence quotient, extra digits on his feet and  
8 fused digits in one of his hand. He had widely spaced eyes, depressed nasal bridge, and high-pitched sound.  
9 He also had a small penis, a small and soft testis and a redundant scrotum. Ultrasound of the abdomen and  
10 pelvis showed mild right-side hydronephrosis. His BMI was 28 and HbA1c was 18.3% and values of LH and  
11 FSH were decreased. He was diagnosed with a case of BBS. He was commenced on insulin for the treatment  
12 of his high blood sugar levels.

13 The initial clinical presentation of BBS may vary from person to person. Thorough examination and a high  
14 degree of clinical suspicion is required to diagnose it. Its management depends upon the signs and symptoms  
15 of the patient. Various options including weight loss, exercise, oral hypoglycemic drugs, and insulin are  
16 available to treat diabetes mellitus in BBS. Some cases can have very high levels of HbA1c on initial  
17 presentation, which may require insulin without trying oral hypoglycemic drugs to treat diabetes mellitus in  
18 BBS.  
19

Accepted, in press

1 **FIGURES AND TABLES.**

2

3 **Figure 1:** Bilateral Polydactyly.



4

5

6

7

8

**Figure :** Syndactyly in Right Hand.



9

10

11

12

13

14

15

16

17

18

19

20

1 **Table 1:** Laboratory Investigations  
2

Name of Investigation	Result	Normal range	Unit
Sodium	132	135-150	mmol/L
Potassium	3.06	3.5-5.1	mmol/L
Chloride	77.8	96-112	mmol/L
Blood Urea	35	18-45	mg/dl
Creatinine	0.4	0.42-1.06	mg/dl
Total Bilirubin	0.3	0.1-1.0	mg/dl
Alanine Transaminase	13	10-50	IU/L
Alkaline Phosphatase	120	35-104	IU/L
Uric Acid	4.7	3.4-7.0	mg/dl
Glucose (Random)	360	70-140	mg/dl
T-3, Total	1.26	0.8-2.0	ng/ml
T-4, (Thyroxin)	11.48	5.1-14.1	ug/dl
Thyroid Stimulating Hormone	0.722	0.3-4.2	uIU/ml
Luteinizing Hormone (LH)	0.01	0.57-12.07	IU/L
Follicle Stimulating Hormone (FSH)	0.79	0.95-11.95	IU/L
Serum Testosterone	17.48	7-75	ng/dl for 12-16 years of age
White Blood Cells	10.5	4-11	X10 <sup>3</sup> /ul
Red blood Cells	4.55	4-6	X10 <sup>3</sup> /ul
Haemoglobin	11.9	11.5-17.5	g/dl
HbA1C	18.3%	4.5-7	%
Serum Calcium	9.2	8.5-10.5	mg/dl
Virology			
Hepatitis B surface Aantign(By ICT)	Negative		
Anti Heatitis C Virus (By ICT)	Negative		
Anti Human immunodeficiency Virus (By ICT)	Negative		

3  
4 mmol/L: millimoles per litre; mg/dl: milligrams per decilitre; ng/ml: nanograms per millilitre; g/dl: grams per  
5 deciliter; IU/L: international units per litre; ug/dl: micrograms per decilitre; U/L: Units per litre; uIU/ml: micro-  
6 international units per milliliter; ng/dl: nanograms per deciliter; ul: microlitre  
7  
8

Accepted, in-press