

Correspondence: Yosuke Kakisaka.
Address: Department of Epileptology, Tohoku University Graduate School of Medicine, 2-1 Seiryō-machi, Aoba-ku, Sendai, Miyagi 980-8575, Japan.
Email: kakisuke@mui.biglobe.ne.jp

References

1. Kakisaka Y, Fujikawa M, Gaillard S. Writing case reports: teaching and tuition techniques, and the improvement of clinical diagnostic reasoning. *Int J Med Students*. 2016 Sep-Dec;4(3):88-90.
2. Niell LS. Writing case reports for the clinical literature: practical approach for the novice author. *J Grad Med Educ*. 2011 Sep;3(3):445.
3. Juyal D, Thaledi S, Thawani V. Writing patient case reports for publication. *Educ Health (Abingdon)*. 2013 May-Aug;26(2):126-9.
4. Kakisaka Y, Hino-Fukuyo N, Kobayashi T, Kubota Y, Kikuchi A, Endo W, et al. Hallucinations associated with cerebrospinal fluid leakage after a lumbar puncture. *Br J Anaesth*. 2012 Sep;109(3):465-6; author reply 466.
5. Loures V, Savoldelli GL, Alberque C, Haller G. Post-dural puncture cerebrospinal fluid leak presenting as an acute psychiatric illness. *Br J Anaesth*. 2012 Mar;108(3):529-30.

Acknowledgments: None.

Conflict of Interest Statement & Funding: The authors have no funding, financial relationships, or conflicts of interest to disclose.

Author Contributions: Conceptualization: YK. Writing, Critical revision of the manuscript, Approval of the final version: YK, MF, SG.

Cite as: Kakisaka Y, Fujikawa M, Gaillard S. Structured didactic education program for writing case reports can motivate medical students. *Int J Med Students*. 2016 Sep-Dec;4(3):131-2.

Latin America: Should We Be Worried about Zika Virus?

About the Author: *Joselyn Ye is currently a fifth-year medical student at Cayetano Heredia University, Lima, Peru of a seven-year program.*

To the Editor,

The Zika virus (ZIKV) was identified for the first time in 1947, almost 70 years ago. Despite being known during all these decades, it has been neglected until the current outbreak in Brazil. This an emergent disease that has been developing slowly throughout these years; we still have limited information and scientific knowledge about this illness and its potential outcomes.¹

This infection causes fairly mild effects, as 80% of the patients go unnoticed.² The most common symptoms are fever, rash, joint pain, and conjunctivitis, which last for several days to a week after the infection.³ The primary mode of transmission is through the bite of an infected mosquito of the *Aedes* species. Apart from mosquitoes, other non-vector means of transmission are sexual intercourse and during pregnancy. Furthermore, there is a strong possibility that ZIKV can be spread through blood transfusions. It is suggested that breastfeeding could be another way of transmission, due to the fact that it has been found in breast milk. However, there are no reports up to date about other means of transmission. Because of the benefits of breastfeeding, mothers are encouraged to breastfeed even in areas where ZIKV is found (Available from: <http://www.cdc.gov/zika/about/overview.html>; cited 2016 Jul 28).

Since ZIKV can cross the blood brain barrier,⁴ a relationship with some severe congenital neurologic defects, such as mi-

crocephaly and Guillian-Barré syndrome, has been established. Mlakar et al described that the cases of microcephaly increased by a factor of 20 among newborns in the northeast of Brazil.^{4,5} Many other authors also describe severe brain injuries, inflammatory signs and calcifications, leading them to think that it is likely that an intrauterine infection brings about the abnormal brain development, and can lead to cerebral palsy after birth.¹⁻³ It is also suggested that the low maternal supplementation of nutrients may affect other organs as well,⁶ although this is yet to be determined.

The World Health Organization declared ZIKV a ‘Public Health Emergency of International Concern’;⁷ as it threatens to become a global pandemic. Up to date, there is no cure or vaccine available (Available from: <http://www.cdc.gov/zika/about/overview.html>; cited 2016 Jul 28). There are different methods for diagnosing ZIKV. One of them is the real-time reverse transcription-polymerase chain reaction (rRT-PCR), which must only be conducted on urine or serum samples collected during the first 14 days after symptoms onset. This test has a high sensitivity and a poor specificity. If the rRT-PCR is negative in a high risk patient, the ELISA for qualitative detection of IgM (Zika MAC-ELISA) is used.⁸ However, a cross-reaction with other flaviviruses is present, as such the test has poor specificity in people living in endemic areas that have been previously exposed to the dengue and yellow fever viruses.⁶⁻⁹ Therefore, results may be difficult to interpret in Latin America.

System-wide barriers impede health care delivery in our region. Latin American health providers have the obligation to generate awareness amongst our people about ZIKV. The need for a stronger surveillance system, suitable diagnostics methods, communication about ZIKV outbreak risks, vector control and adequate guidance to pregnant women is necessary.³

Joselyn Ye,¹ Marcelo Bedoya.¹

¹ Faculty of Medicine, Cayetano Heredia University, Lima, Peru.

Correspondence: Joselyn Ye.
Address: Facultad de Medicina Alberto Hurtado. Universidad Peruana Cayetano Heredia, Lima, Perú.
Email: joselyn.ye.t@upch.pe

References

1. Schuler-Faccini L, Ribeiro EM, Feitosa IM, Horovitz DD, Cavalcanti DP, Pessoa A, et al. Possible association between Zika virus infection and microcephaly – Brazil 2015. *MMWR Morb Mortal Wkly Rep*. 2016 Jan;65(3):59-62.
2. Petersen E, Wilson ME, Touch S, McCloskey B, Mwaba P, Bates M, et al. Rapid spread of Zika virus in the Americas – implication for public health preparedness for mass gatherings at the 2016 Brazil Olympic Games. *Int J Infect Dis*. 2016 Mar;44:11-5.
3. Heymann DL, Hodgson A, Sall AA, Freedman DO, Staples JE, Althabe F, et al. Zika virus and microcephaly: why is this situation a PHEIC? *Lancet*. 2016 Feb 20;387(10020):719-21.
4. Tetro JA. Zika and microcephaly: causation, correlation, or coincidence? *Microbes Infect*. 2016 Mar;18(3):167-8.
5. Mlakar J, Korva M, Tul N, Popovic M, Poljsak-Prijatelj M, Mraz J, et al. Zika virus associated with microcephaly. *N Engl J Med*. 2016 Mar 10;374(10):951-8.
6. World Health Organization. WHO statement on the first meeting of the International Health Regulations. Emergency Committee on Zika virus and observed increase in neurological disorders and neonatal malformations; 2016.
7. Vogel G. Evidence grows for Zika virus as pregnancy danger. *Science*. 2016

Submission: Sep 17, 2016
Acceptance: Nov 14, 2016
Publication: Dec 24, 2016

Mar 11;351(6278):1123-4.

8. Lazear HM, Stringer EM, de Silva AM. The emerging Zika virus epidemic in the Americas: research priorities. *JAMA*. 2016 May 10;315(18):1945-6.

9. Narayanan R. Zika virus therapeutics: drug targets and repurposing medicine from the human genome. *MOJ Proteomics Bioinform*. 2016;3(3):00084.

Acknowledgments: The authors would like to express their gratitude to everyone who supported them throughout their undergraduate education, especially Instituto de Medicina Tropical Alexander von Humboldt, which provided them with the knowledge and guidance to write this manuscript.

Conflict of Interest Statement & Funding: The authors have no funding, financial relationships, or conflicts of interest to disclose.

Author Contributions: Conceptualization, Data collection, Writing, Critical revision of the manuscript: JY, MB. Approval of the final version: JY.

Cite as: Ye J, Bedoya M. Latin America: should we be worried about Zika virus? *Int J Med Students*. 2016 Sep-Dec;4(3):132-3.

Methylprednisolone-Responsive Leptospiral Acute Pulmonary Syndrome

About the Author: Professor Santos is a Professor-adjunct of Internal Medicine at Armed Forces Hospital and Catholic University of Brasília, Brasília-DF, Brazil. He has half a century of docent activities.

To the Editor,

The article by Johan-Arief et al highlights the role of early methylprednisolone therapy to reduce morbidity and mortality of the acute pulmonary syndrome in leptospirosis,¹ and the authors emphasize the ideal administration of this drug within 12 hours after diagnosis. They reported a 26-year-old man with fever of one-week duration, who on the third day had presented with cough and breathlessness, and episodes of hemoptysis occurred two days later.¹ His professional work involved frequent close contact with soil, and he was swimming in a waterfall a week before the onset of the high-grade fever associated with chills, rigors and body aches.¹ Physical signs were jaundice, subconjunctival suffusion, and bilateral pulmonary crepitations. Blood tests showed neutrophilic leucocytosis, low platelets, disordered coagulation, liver and renal functions; and the chest x-ray revealed scattered alveolar infiltrates. The Weil's disease was diagnosed by positive IgM ELISA, and the specific microscopic agglutination test (1:400).¹ Intravenously, he underwent tranexamic acid (500 mg 3 times daily), plus benzyl penicillin (1200 mg every 6 hours) during a week, and methylprednisolone (15 mg/kg daily) for 3 days.¹ His clinical response to treatment was good and he was soon discharged home asymptomatic. Environmental factors and typical features of severe leptospirosis raised the diagnostic suspicion; as the early diagnosis is the cornerstone of prompt adequate treatment of severe leptospirosis, one must consider some main concerns that constitute frequent challenges in clinical practice.^{2,3} During the first week of acute leptospirosis, the blood borne bacteria almost invariably do not evoke an immune response enough to yield antibody levels and false negative tests can occur;^{1,3} so, the goal of starting adequate therapy within the first seven days of disease may be hindered.³ Pulmonary involvement with hemoptysis has been more often reported in severe Weil's disease, and the coexistence of accentuated thrombocytopenia may play a role on the poorest prognoses. Significant pulmonary changes as well as low platelet counts can pose other concerns about diagnostic pitfalls with Hantavirus and dengue

infections, especially in developing countries.²⁻⁵ Since the first reports in North American Indians, the possibility of mistakes involving the acute pulmonary syndrome caused by leptospirosis and HPS was commented in Brazilian Journals.⁴ Brazilian authors described a 19-year-old man with the diagnosis of anicteric leptospirosis, who had a productive cough without hemoptysis, reduced lung sounds on the right lower third of the thorax, and normal kidney function. Images of computed tomography confirmed the acute inflammatory process in the right lung, and he underwent a successful course of intravenous ceftriaxone.³ Worthy of note, the patient had clinical and electrocardiographic findings of pericarditis; moreover, the count of bands was high (up to 1020) and the platelet count was low ($82 \times 10^9/L$). Although not confirmed, these clinical and complementary findings were suggestive of HPS; a condition that has been described in the Brazilian Central Plateau either isolated or as a coinfection.^{3,5} On a practical standpoint, corticosteroids have been useful to control both the syndromes.^{1,5}

Vitorino Modesto Santos.¹

¹ Armed Forces Hospital and Catholic University, Brasília-DF, Brazil.

Correspondence: Vitorino Modesto Santos.

Address: Armed Forces Hospital and Catholic University, Brasília-DF, Brazil.

Email: vitorinomodesto@gmail.com

References

- Johan-Arief ID, Lee SH, Er XY, Kasinathan G, Pillai N. Methylprednisolone-responsive leptospiral acute pulmonary syndrome: a case report. *Int J Med Students*. 2015 Sep-Dec;3(3):159-62.
- Dos Santos VM. Manifestations and complications of leptospirosis. *Med J Islam Repub Iran*. 2016 Mar 2;30:337.
- Santos VM, Santos UM, Gebrin DG, Santos AM, Cancado AC. Anicteric leptospirosis with pneumonitis, pericarditis and acalculous cholecystitis. *Infez Med*. 2014 Sep;22(3):236-40.
- dos Santos VM. [Leptospirosis and pulmonary hantavirus syndrome]. *Rev Assoc Med Bras* (1992). 1994 Jul-Sep;40(3):225. Portuguese
- Pinto Junior VL, Hamidad AM, Albuquerque Filho Dde O, dos Santos VM. Twenty years of hantavirus pulmonary syndrome in Brazil: a review of epidemiological and clinical aspects. *J Infect Dev Ctries*. 2014 Feb;8(2):137-42.

Acknowledgments: None.

Conflict of Interest Statement & Funding: The author has no funding, financial relationships, or conflicts of interest to disclose.

Author Contributions: Conceptualization, Writing: VMS.

Cite as: Santos VM. Methylprednisolone-responsive leptospiral acute pulmonary syndrome. *Int J Med Students*. 2016 Sep-Dec;4(3):133.

Submission: Sep 26, 2016

Acceptance: Dec 07, 2016

Publication: Dec 25, 2016