

Fulminant Hepatic Failure as the Initial Presentation of Hodgkin's Disease and Liver Transplantation: A Case Report

Mauricio Alejandro Saldaña-Ruiz,¹ Federico Ortiz-Alonso,² Adriana Carolina Sandoval-González,² Liliana Sayuri Tapia-Brito,² Laura Carolina Lozano-Galván,³ Karla Monserrat Ramírez-Pintor.⁴

Abstract

Background: Hodgkin's disease, a B-cell neoplasm, primarily impacts lymph nodes or extranodal lymphoid tissue. It includes two distinct entities: classical (95%) and lymphocyte-predominant nodular. While the disease commonly manifests as the growth of cervical and intrathoracic lymph nodes in 60-90% of cases, there are rare instances where Hodgkin's disease has been linked to fulminant liver failure, carrying a very poor prognosis. **The Case:** We present the case of a 13-year-old Hispanic female, who started with an insidious condition that evolved to fulminant hepatic failure of unknown etiology with an AST of 770 mg/dl. It was decided to perform an orthotopic liver transplant, the histopathological analysis of the explant and a lymph node reported mixed cellularity Hodgkin's disease. Subsequently, the hematology service requested a lumbar puncture, with no evidence of infiltration. It was decided to initiate six cycles of chemotherapy (CTX) with BEACOPP (bleomycin, etoposide, adriamycin, cyclophosphamide, vincristine, procarbazine, and prednisone) scheme, evolving without complications and achieving a complete response eleven months later; currently, she has been free of disease for three years. **Conclusion:** The etiology of Hodgkin's disease in our 13-year-old patient remains elusive, emphasizing the importance of early diagnosis and diverse treatment approaches. Despite limited hospital resources, the decision to proceed with the transplant was driven by the potentially fatal outcome if left untreated. Future considerations may necessitate individualizing each case, and carefully assessing the risks and benefits associated with transplantation.

Introduction

Hodgkin's disease is a neoplasm of B lymphocytes, that affects the lymph nodes and has a bimodal distribution, with the first peak between 15 and 30 years of age and the second around 55 years of age.¹ Moreover, the World Health Organization classifies this neoplasm into two types: predominantly lymphocytic nodular Hodgkin's disease (5%) and classical Hodgkin's disease with the presence of Reed-Sternberg cells (95%).²

It affects different parts of the body, leading to variations in its clinical presentation. The most common presentation involves cervical and intrathoracic lymph node growths in 60-80%,³ contrasting with the rare initial presentation through hepatic infiltration, since it constitutes approximately 0.44% of all cases,⁴ the clinical presentation is usually nonspecific, leaning more towards one consistent with fulminant hepatic failure, which is defined as the development of coagulopathy and encephalopathy within eight weeks of the onset of hepatic dysfunction in patients without the pre-existing liver disease,⁵ many of these cases with fatal outcomes, however, for years it has been an absolute contraindication to receive a transplant to have an untreated active neoplasm,⁶ and it is still maintained to this day.⁷ We present a case of a 13-year-old female who received a liver transplant at the time she had Hodgkin's disease.

Highlights:

- Currently, receiving a liver transplant having a neoplasm is a contraindication.
- The presentation of Hodgkin's disease with hepatic infiltration is extremely rare.
- There are only a few reported cases of liver transplantation at the time the patient has a neoplasm.

The Case

A 13-year-old Hispanic female, previously healthy, who began 6 weeks earlier with fatigue, nausea, and fever, went to the doctor for evaluation and requesting laboratory tests, iron deficiency anemia was diagnosed, being managed with acetaminophen at adequate doses for her weight and oral supplemental iron at conventional doses, remaining asymptomatic for six weeks. Afterwards she restarted with asthenia, intermittent fever, nausea and vomiting, going again for medical evaluation, finding data of pancytopenia, jaundice, abdominal pain, nausea, vomiting, hyperlipidemia and hyperthermia for which she was sent to the local emergency department, on admission in poor general condition, with evidence of grade I encephalopathy, generalized icteric tinge, hepatosplenomegaly, paraclinical data included

¹ Social Service. Instituto Mexicano del Seguro Social, Monterrey, México.

² MD. Instituto Mexicano del Seguro Social, Monterrey, México.

³ MD. Facultad de Medicina UANL, Monterrey, México.

⁴ Sixth-year Medical Student. Facultad de Medicina UANL, Monterrey, México.

About the Author: Mauricio Saldaña is currently a medical student of Facultad de Medicina UANL, Monterrey, Mexico doing his social service at Instituto Mexicano del Seguro Social in the Coordination of Donation and Transplants. He presented a poster at the 3rd Medical Student World Congress of Neurosurgery on September 1st in Germany.

Correspondence:

Mauricio A. Saldaña-Ruiz.

Address: Moderna, 27170 Torreón, Coahuila, Mexico.

Email: mauriciomgr14@gmail.com

Editor: Francisco J. Bonilla-Escobar
Student Editors: Rebecca Murerwa,
Andrea Cuschieri & Esther Bassey
Copyeditor: Eugenia M. Ramos-Dávila

Submission: Oct 18, 2023
Revisions: Nov 30, 2023
Responses: Dec 9, 2023
Acceptance: Dec 11, 2023
Publication: Dec 11, 2023
Process: Peer-reviewed

anemia, transaminase elevation, cholestasis, alterations in coagulation times test and findings suggestive of acute liver failure (Table 1). Supportive management was initiated, and the patient was admitted to the pediatric intensive care unit, broadening the diagnostic approach to determine the etiology. Upon questioning family members, there was no reported ingestion of herbal medicine, illicit substances, or others.

She was evaluated by the transplant service of the hospital (Table 2), fulfilling the criteria for liver transplantation, classifying the case as UNOS status 1A. The next day, a cadaveric liver transplant was performed, and the explant with a lymph node was sent to pathology for histopathological studies. The report indicated a liver weight of 1.160 kg, measures 21x17x5cm, irregular surface, rough, greenish color with purplish areas. The lymph node measures 1.1x0.6cm, grayish-brown, irregular, and soft. The pathology report shows a hepatic hilum lymph node consistent with Hodgkin's disease of mixed cellularity and hepatic infiltration; immunohistochemistry is positive for CD15, CD30, CD45, and fascin in neoplastic cells. A week later, a bone marrow biopsy was performed, without evidence of neoplastic infiltration. Afterwards, the Hematology service classified the lymphoma as stage IV.

Due to the diagnosis, management with chemotherapy BEACOPP (Bleomycin, Etoposide, Adriamycin, Cyclophosphamide, Vincristine, Procarbazine and Prednisone) scheme of six cycles was started, which lasted for nine months; during her management a cervical lymph node was found in Computed Tomography, which was kept under follow-up, and a biopsy of the lymph node was performed without finding infiltration. The rest of the clinical evolution was towards improvement, maintaining stable liver enzyme levels throughout the CTX sessions, complying with immunosuppressive management, achieving complete response, currently free of disease for three years and in her last consultation with good general condition and adequate hydration, as well as steady laboratory levels (Table 3).

Discussion

It can be seen in the case as the patient evolved satisfactorily achieving a complete response and remaining free of disease at present, even though different guidelines mention that it is a contraindication to perform a liver transplant in fulminant hepatic failure secondary to neoplasms,⁷ some cases have been reported in the literature of favorable evolutions after undergoing liver transplantation.

Kirsten M. et al. reported a case involving a seven-year-old male with fulminant hepatic failure requiring emergency liver transplantation, during surgery, enlarged lymph nodes were located around the portal vein, and Hodgkin's disease was later diagnosed, the patient had a relapse when an enlarged cervical lymph node was located, so chemotherapy was performed, achieving a complete response, even some time later he needed a retransplantation at 13 months due to chronic graft rejection.⁸

Table 1. Blood Test Results After Admission.

Paraclinical Report	Value	Normal Values
Hemoglobin	10.6	12.2-18.1 g/dL
Leukocytes	3.8	4.2-10.2 K/uL
Neutrophils	2.4	2.0-6.9 K/uL
Lymphocytes	0.1	0.6-3.4 K/uL
Platelets	94.7	142-424 K/uL
Hepatitis B Virus (HBV)	Negative	Not applicable
Hepatitis C Virus (HCV)	Negative	Not applicable
Total bilirubin	12.04	0.1-1.2 mg/dl
Direct bilirubin	8.86	<0.3 mg/dl
Indirect bilirubin	3.18	0.2-1.2 mg/dl
Alanine Aminotransferase (ALT)	1144	4-36 U/L
Aspartate Aminotransferase (AST)	770	8-33 U/L
Human Immunodeficiency Virus (HIV)	Negative	Not applicable
Lupus anticoagulant	Normal	<1.2

Table 2. Blood Test Results to Evaluation for Transplantation Department.

Paraclinical Report	Value	Normal Values
Total bilirubin	17	0.1-1.2 mg/dl
Direct bilirubin	11.9	<0.3 mg/dl
Indirect bilirubin	5.6	0.2-1.2 mg/dl
ALT	1014	4-36 U/L
AST	716	8-33 U/L
Lactate Dehydrogenase (LDH)	418	105-333 U/L
International Normalized Ratio (INR)	8.3	1
Ammonium	285	15-45 µg/dl
Procalcitonin	>10	<0.5 ng/ml

Table 3. Last Consult Blood Test Results.

Paraclinical Report	Value	Normal Values
Hemoglobin	13	12.2-18.1 g/dl
Leukocytes	3.9	4.2-10.2 K/uL
Platelets	126	142-424 K/uL
INR	1	1
Total bilirubin	0.30	0.1-1.2 mg/dl
ALT	19	4-36 U/L
AST	24	8-33 U/L
Sirolimus levels	6.8	5-15 ng/ml

Brannigan et al. reported the case of a 12-year-old pediatric patient who developed fulminant hepatic failure, initially, parvovirus was considered as the main cause, a living-donor liver transplant was performed with his mother; when the explant was examined, Hodgkin's disease was diagnosed, the patient up to the date of publication remains in remission at one year.⁹

There are other cases in which a liver transplant was performed despite a previous diagnosis of Hodgkin's disease, which is also considered a contraindication at present.

Frank S Hong et al. reported two cases of fulminant hepatic failure, one of a 19-year-old male with a pre-transplant diagnosis of Hodgkin's disease who achieved a complete response to the disease; the other 55-year-old patient died without having received a transplant due to posterior fossa hemorrhage.¹⁰

Also, Hope et al. reported a similar case of a five-year-old pediatric patient with a history of Hodgkin's disease diagnosed at the age of two, who developed fulminant hepatic failure, a percutaneous biopsy was performed in which areas of portal infiltration by lymphocytes were found without detection of cells suggesting cancer, It was decided to perform the transplant, during surgery a plaque was located in the patient's liver, and an intraoperative biopsy was performed with a high possibility of being Hodgkin's disease, it was decided to continue with the transplant as it was not entirely diagnostic; histopathologic examination was performed diagnosing Hodgkin's disease, sometime later there was a recurrence of cancer, which was successfully treated with rituximab, the patient is in remission five years later with no apparent complications.¹¹

Although it is not intended that all patients with hepatic infiltration by undiagnosed Hodgkin's disease who evolve to fulminant hepatic failure receive a transplant, due to the scant information available to date, it is important to observe how the

reported cases have achieved favorable results, likely since this type of neoplasm has good survival rates depending on the stage.

We believe it would not be appropriate to dismiss this hypothesis. Therefore, we might consider, in the future, individualizing each case to evaluate whether receiving the transplant would be beneficial or not.

In addition, Given the limited resources in many hospitals to obtain an early diagnosis meeting the criteria for Hodgkin's disease, it was decided to perform the transplant since, it had not been performed, it is very likely that the patient would have a fatal outcome.

Summary – Accelerating Translation

Falla hepática fulminante como la presentación inicial de enfermedad de Hodgkin y trasplante hepático: Reporte de un caso.

Desde hace tiempo es una contraindicación recibir un trasplante al tener una neoplasia diagnosticada y se mantiene hasta nuestros días, sin embargo, ha habido casos reportados en la literatura sobre pacientes que recibieron un trasplante hepático en esas mismas circunstancias. En este caso presentamos a una paciente femenina la cual inicio con un cuadro insidioso, posteriormente progreso a falla hepática fulminante de etiología desconocida, decidiéndose realizar un trasplante hepático de receptor cadavérico no emparentado al cumplir con criterios del King's College. Posteriormente se reportó enfermedad de Hodgkin en el hígado y un ganglio peri hepático, iniciándose quimioterapia con esquema BEACOPP (bleomicina, etopósido, adriamicina, ciclofosfamida, vincristina, procarbazona, y prednisona), evolucionando favorablemente y estando libre de enfermedad en la actualidad.

Es importante ver como la paciente evoluciona de manera favorable y que, en el caso que se hubiera diagnosticado primero la neoplasia, no hubiera recibido el trasplante y probablemente hubiera fallecido por complicaciones mismas de la falla hepática fulminante. Por lo cual pudiera ser de importancia a futuro para individualizar cada caso y decidir si se beneficiaría de recibir un trasplante o no.

References

- Smith A, Howell D, Patmore R, Jack A, Roman E. Incidence of haematological malignancy by sub-type: a report from the Haematological Malignancy Research Network. *Br J Cancer* 2011;105:1684-92.
- Siegel RL, Miller KD, Jemal A. Cancer statistics, 2017. *CA Cancer J Clin* 2017;67(1):7-30.
- Jiménez-Ríos G, Balderas-Delgado C, Vargas-Bravo CA, Díaz-Hernández PI, Velasco-Vargas CA, Velázquez-Guerrero D et al. Experience of haematopoietic progenitor cell transplantation in Hodgkin's lymphoma at the Regional High Speciality Hospital of Ixtapaluca. *Rev Mex Traspl*. 2023; 12 (1): 45-9. Spa
- Karmacharya P, Bhandari N, Aryal MR, Pandit AA, Pathak R, Ghimire S et al. Before it crumbles: Fulminant Hepatic Failure secondary to Hodgkin's Lymphoma. *J Community Hosp Intern Med Perspect*. 2014; 4 (5): 25821.
- Trey C, Davidson CS. The management of fulminant hepatic failure. *Prog Liver Dis* 1970; 3: 282-98
- Rowbotham D, Wendon J, Williams R. Acute liver failure secondary to hepatic infiltration: a single centre experience of 18 cases. *Gut*.1998;42:576-80.
- Squires JE, Alonso EM, Ibrahim SH, Kasper V, Kehar M, Martinez M et al. North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition Position Paper on the Diagnosis and Management of Pediatric Acute Liver Failure. *J Pediatr Gastroenterol Nutr*. 2022; 74(1):138-58.
- Woolf KMW, Wei MC, Link MP, Arber DA, Warnke RA. Nodular lymphocyte-predominant Hodgkin lymphoma presenting as fulminant hepatic failure in a pediatric patient: a case report with pathologic, immunophenotypic, and molecular findings. *Appl Immunohistochem Mol Morphol*. 2008;16(2):196-201.
- Brannigan L, Etheredge HR, Beretta M, Demopoulos D, Bennett KG, Beringer N et al. Successful ABO-incompatible living donor liver transplant for acute liver failure secondary to Hodgkin's Lymphoma in a child. *Pediatr Transplant*. 2020; 24(7):e13796.
- Hong FS, Smith CL, Angus PW, Crowley P, Ho WK. Hodgkin lymphoma and fulminant hepatic failure. *Leuk Lymphoma*. 2010; 51(5):947-51.
- Hope BC, Chau KY, Evans HM, Mout S, Munn S, Yeong ML et al. Hodgkin disease relapse discovered at the time of liver transplant for acute liver failure. *Pediatr Transplant*. 2012; 16(1): E10-E14.

Acknowledgments

The authors acknowledge the support of José Guillermo Martínez Flores and Cindy Sharon Ortiz Arce for their help with manuscript writing.

Conflict of Interest Statement & Funding

The Authors have no funding, financial relationships or conflicts of interest to disclose.

Author Contributions

Conceptualization: MASR, and LCLG. Validation: MASR, FOA, ACGS, and LSTB. Formal Analysis: MASR. Investigation: MASR, LCLG, and KMRP. Resources: MASR, LCLG, and KMRP. Writing – Original Draft: MASR, LSTB, and LCLG. Writing – Review & Editing: MASR, ACGS, LSTB, and KMRP. Visualization: MASR, and KMRP. Supervision: MASR, and FOA. Project Administration: MASR.

Cite as

Saldaña-Ruiz MA, Ortiz-Alonso F, Sandoval-González AC, Tapia-Brito LS, Lozano-Galván LC, Ramírez-Pintor KM. Fulminant Hepatic Failure as the Initial Presentation of Hodgkin's Disease and Liver Transplantation: A Case Report. Int J Med Stud. 2024 Jan-Mar;12(1):92-95.

This work is licensed under a [Creative Commons Attribution 4.0 International License](#)

ISSN 2076-6327

This journal is published by [Pitt Open Library Publishing](#)

