

ORIGINAL RESERCH

50. Healthcare Access Dimensions among ESRD Patients on Dialysis in Northern Sudan during Wartime 2024 Ivas Abdelhadi¹

¹University of Khartoum, Faculty of Medicine, Sudan

Background: Patients with end-stage renal disease (ESRD) undergoing hemodialysis are among the most vulnerable groups in low-income and conflict-affected settings. Adequate access to healthcare is crucial for maintaining treatment continuity, preventing complications, and supporting nutritional health in this population. This study assesses healthcare access and its determinants among end-stage renal disease (ESRD) patients on hemodialysis in Northern Sudan and examines the relationship between access and malnutrition.

Methods: This cross-sectional study involved secondary data analysis from 137 ESRD patients in four dialysis centers in Northern Sudan using cluster sampling. Healthcare access was measured using a structured questionnaire based on the modified Penchansky and Thomas model. Nutritional status was assessed using the Subjective Global Assessment (SGA). Descriptive statistics, bivariate analysis, and logistic regression were used to identify determinants and predictors of healthcare access using the Statistical Package for Social Science (SPSS) v21. Statistical significance was set at p-value < 0.05.

Results: Overall, all patients showed moderate to high levels of healthcare access (mean index: 3.87 ± 0.51). Affordability scored the lowest among access dimensions. Higher income and age were significant predictors of better access (AOR=9.99 and AOR=3.87, respectively; p<0.05), while gender, displacement, and health insurance showed no significant contribution. Malnourished patients showed significantly lower scores on the affordability and awareness domains (p-value < 0.05).

Conclusion: While patients in Northern Sudan had moderate to high access to healthcare, affordability was the most prominent barrier, being more prominent in malnourished patients, besides the low awareness. We recommend socioeconomic support, facilitating healthcare access for vulnerable patients, and nutritional educational programs.

Table 1. Baseline characteristics and healthcare access and use of the ESRD on Haemodialysis in Northern Sudan. N=141

| Sociodemographic and the Patients | clinical characteristics of | N (%) |
|---------------------------------------|---|------------------------|
| rudend | Karima Teaching | 54 (38.3) |
| | Hospital | ` ' |
| Treating Centre | Algolid Dialysis Centre Argo Dialysis Centre | 44 (31.2) 24 (17.0) |
| | Wadi Halfa Dialysis | 19 (13.5) |
| | Centre < 45 | 44 (31.2) |
| Age | 45-65 | 66 (46.8) |
| | >= 65 | 31 (22.0) |
| | M(Sd) Male | 52 (14.5) |
| Gender | Female | 96 (68.1) 45 (31.9) |
| Marital Status Education | Single | 22 (15.6) |
| | Married Divorced or widowed | 105 (74.5) |
| | Missing | 13 (9.2) 1 (0.7) |
| | Illiterate | 12 (8.5) |
| | Primary | 49 (34.8) |
| | Secondary University or higher | 48 (34.0) 31 (22.0) |
| | Missing | 1 (0.7) |
| Occupation | Retired or non-working | 111 (78.7) |
| | Free worker Formal Employee | 19 (13.5) 10 (7.1) |
| | Missing | 1 (0.7) |
| | Less than 150K | 80 (56.7) |
| Income/ Month in SDG | 150-300 Above 300K | 41 (29.1) |
| | Missing | 18 (12.8) 2 (1.4) |
| Displacements | Yes | 70 (50.0) |
| | No No | 70 (50.0) |
| | Missing Own house | 1 (0.7) 48 (34.0) |
| Household | Rented house | 11 (7.8) |
| | Relative/extended | 78 (55.3) |
| | family IDP camp | 3 (2.1) |
| | Missing | 1 (0.7) |
| Residence | Rural | 89 (63.1) |
| | Urban Missing | 47 (33.3) 5 (3.5) |
| | No | 12 (8.5) |
| | Missing | 1 (0.7) |
| | Less than 2 years | 39 (27.7) |
| Haemodialysis Duration | 2-5 years More than 5 years | 44 (31.2) 58 (41.1) |
| | Mean (SD) | 4.6 (4.7) |
| Health care access | Low Level | 0 (0) |
| | Moderate Level High Level | 54 (38.3) 83 (58.9) |
| | Missing | 4 (2.8) |
| | Mean Access Index (SD) | 3.9 (0.5) |
| | SHI Private | 99 (70.2) 6 (4.3) |
| Health Insurance | Non | 34 (24.1) |
| | Missing | 1 (0.7) |
| Outpatient Healthcare | Non 1-2 | 73 (51.8) 39 (27.7) |
| service contacts (6- months) | 3 or more | 15 (10.6) |
| | Missing | 4 (2.8) |
| Where do you head when you feel sick? | PHC centres Traditional healer | 12 (8.5) |
| | ER /GP in a 2ndary | 12 (8.5) 89 (63.1) |
| | facility | 03 (05.1) |
| | Specialized care | 23 (16.3) |
| Travel time to facility | Less than 30 mins 30 mins – 1 hr | 73 (51.8) 49 (34.7) |
| maver time to facility | More than 1 hr | 19 (13.5) |

This work is licensed under a <u>Creative Commons</u>
<u>Attribution 4.0 International License</u>

ISSN 2076-6327

This journal is published by Pitt Open Library
Publishing

