

1 **Title:** Occupational Exposure to Blood and Body Fluids and Its Association with Anxiety Among Final-Year
2 Medical Students: A Single-Center Cross-Sectional Study

3
4 **Article type:** Original Article

5
6 **Author names:**

- 7 1. Levent ÖZDEMİR
- 8 2. Sena Sude ÖZYAR
- 9 3. Sabrina İSRAFİLZADE
- 10 4. Behrokh KHORASANI

11
12 **Degrees and Affiliations:**

- 13 1. Prof. Dr., MD. Bursa Uludağ University, Faculty of Medicine, Public Health Department, Bursa/TURKEY
- 14 2. 6th year Medical Student, Bursa Uludağ University, Faculty of Medicine, Bursa/TURKEY
- 15 3. 6th year Medical Student, Bursa Uludağ University, Faculty of Medicine, Bursa/TURKEY
- 16 4. 6th year Medical Student, Bursa Uludağ University, Faculty of Medicine, Bursa/TURKEY

17
18 **ORCID (Open Researcher and Contributor Identifier):**

- 19 1. <https://orcid.org/0000-0002-4330-8334>
- 20 2. <https://orcid.org/0009-0009-3891-3618>
- 21 3. <https://orcid.org/0009-0003-4014-7146>
- 22 4. <https://orcid.org/0009-0005-2202-274X>

23
24
25 **About the author:** Sena Sude ÖZYAR is currently a 6th-year medical student at Bursa Uludağ University
26 Faculty of Medicine, Bursa, Turkey, in a six-year program. She has a keen interest in scientific research and
27 aspires to pursue an academic career in the future.

28
29 **Corresponding author email:** leventozdemir@uludag.edu.tr

30 **Acknowledgment:** The authors thank all the final-year medical students who voluntarily participated in this
31 study.

32 **Financing:** None

33 **Conflict of interest statement by authors:** The authors declare that they have no competing interests.
34
35
36
37
38
39
40
41

1 **Authors Contribution Statement:** Conceptualization: LEVENT ÖZDEMİR.
2 Data Curation: SENA SUDE ÖZYAR, SABRINA İSRAFİLZADE, BEHROKH KHORASANI.
3 Formal Analysis: LEVENT ÖZDEMİR, SENA SUDE ÖZYAR.
4 Investigation: LEVENT ÖZDEMİR, SENA SUDE ÖZYAR, SABRINA İSRAFİLZADE, BEHROKH KHORASANI.
5 Methodology: LEVENT ÖZDEMİR.
6 Project Administration: LEVENT ÖZDEMİR.
7 Supervision: LEVENT ÖZDEMİR.
8 Writing - Original Draft: SENA SUDE ÖZYAR.
9 Writing - Review Editing: LEVENT ÖZDEMİR.

10

11

12 **Manuscript word count:** 3287

13 **Abstract word count:** 250

14 **Number of Tables:** 5

15

16 **Personal, Professional, and Institutional Social Network accounts.**

17

- **Facebook:** levent.ozdemir.9678

18

- **Twitter:** @MynameisBehrokh,

19

- **Instagram:** sudeozyar, behrokh.khorasani, @sabrina.izd, lvnt.ozdmr, bursa_uludagtip

20

- **Linkedin:** Sabrina İsrafilzade, Behrokh Khorasani,

21

Accepted, in-press

1 **ABSTRACT**

2 **Background:** Occupational exposure to blood and body fluids (BBFs) remains a significant risk for healthcare
3 professionals, particularly those in training. These exposures not only pose a biological hazard but may also
4 contribute to psychological distress. This study aimed to assess the prevalence of BBF exposure among final-
5 year medical students and investigate its association with anxiety levels.

6 **Methods:** A cross-sectional study was conducted in January–February 2025 at Bursa Uludağ University Faculty
7 of Medicine, Turkey. Of 271 final-year students, 203 participated (74.9%). Data were collected using a structured
8 online questionnaire assessing sociodemographic characteristics, exposure history, and anxiety levels using
9 the Generalized Anxiety Disorder-7 (GAD-7) scale. Predictors of anxiety severity were analyzed using negative
10 binomial regression.

11 **Results:** Overall, 56.2% of students reported at least one BBF exposure, with 67.5% experiencing multiple
12 incidents. Common exposures occurred during venipuncture (50%) and arterial puncture (33.3%), with
13 emergency department rotations posing the highest risk (66.7%). Despite high glove usage (100%), gown usage
14 was low (16.7%). The mean GAD-7 score was significantly higher among exposed students (7.21 vs. 5.39,
15 $p=0.016$). Regression analysis revealed BBF exposure (IRR=1.34), high-risk departments (IRR=1.52), and
16 factors like performance anxiety (IRR=1.85) significantly increased anxiety severity.

17 **Conclusion:** In this single-center study, occupational BBF exposure was highly prevalent among final-year
18 medical students and was significantly associated with elevated anxiety levels, especially in high-pressure
19 clinical settings. Despite existing safety training and orientations, the study's findings revealed persistent gaps
20 in critical areas such as PPE compliance (low gown usage) and effective management of psychosocial stressors
21 (hierarchical pressure).

22
23 **Key Words:** Personal Protective Equipment, medical students, occupational exposure, anxiety, clinical training,
24 curriculum
25

1 INTRODUCTION

2
3 The risk of occupational infection with blood-borne pathogens in healthcare workers is a significant
4 problem worldwide.¹ Each year in Europe, approximately 304,000 healthcare workers sustain percutaneous
5 injuries from materials contaminated with Hepatitis B virus (HBV), 149,000 with Hepatitis C virus (HCV), and
6 22,000 with Human Immunodeficiency Virus (HIV). Notably, 90% of these injuries occur in developing
7 countries.² An estimated 3 million healthcare workers experience percutaneous exposures annually worldwide,
8 with 40% occurring during training.³ Exposure to blood and infectious body fluids represents a significant
9 occupational hazard for healthcare workers and medical students during their clinical training.⁴

10 Sharps injuries, particularly needlestick injuries, impose significant economic burdens globally. Italy
11 reports an average cost of €375 (range: €290-460) per incident for post-exposure management⁵, with 70-80%
12 of occupational exposures involving percutaneous injuries.³ In China, the total economic burden of needlestick
13 injuries among healthcare workers was estimated to be ¥5.8 billion, with approximately half of this cost (¥2.8
14 billion) attributed to nurses.⁶ According to studies conducted in Turkey, approximately 60% of healthcare
15 professionals accidentally come into contact with patients' blood or body fluids at least once.

16 According to studies conducted in Turkey, approximately 60% of healthcare professionals accidentally
17 come into contact with patients' blood or body fluids at least once.⁷⁻⁹

18 In contrast to experienced clinical healthcare providers, medical interns often enter the clinical
19 environment for the first time. Their limited medical experience, combined with an intense eagerness to perform
20 unfamiliar procedures, frequently without sufficient training or supervision, may increase their vulnerability to
21 occupational exposures.¹⁰

22 Anxiety-characterized by persistent worry impairing daily function, can be both a predisposing factor
23 for, and a psychological consequence of, occupational exposures.¹¹ The findings of a meta-analysis suggest
24 that the bidirectional relationship between work injuries and mental health challenges can indeed form a stress-
25 injury cycle.¹²

26 This study aimed to investigate the relationship between BBF exposure and anxiety levels among final-
27 year medical students, and to explore potential implications for medical education and occupational safety
28 training. The insights derived from this investigation are intended to inform the design of medical curricula and
29 educational policies that better prepare students for the realities of clinical practice, focusing on both physical
30 safety and psychological well-being. To achieve this, we employed a cross-sectional design using a validated
31 anxiety screening instrument (GAD-7) and multivariate regression analysis to assess factors associated with

1 anxiety severity. We hypothesized that students who experienced blood or body fluid exposure during clinical
2 training would exhibit significantly higher anxiety scores compared to their unexposed peers.

4 **METHODS**

5 We conducted this cross-sectional study between January and February 2025 among final-year medical
6 students at Bursa Uludağ University Faculty of Medicine. The target population consisted of 271 students. A
7 total of 203 students completed the online questionnaire, yielding a response rate of 74.9%. We did not perform
8 sampling, as the study aimed to reach the entire population. We collected data using a Google Forms survey.
9 The online questionnaire began with an Informed Voluntary Consent Form, and we enrolled participants who
10 agreed to the terms.

11 The questionnaire consisted of two main sections. The first section, consisting of 12 items assessing
12 sociodemographic characteristics and occupational exposure to blood and body fluids, was developed by the
13 research team based on a literature review of similar studies. Questions covered exposure history, routes,
14 associated procedures, clinical rotations, PPE use, and self-reported contributing factors, with multiple
15 selections allowed where applicable. The first section items were not designed as a formal scale, as they were
16 intended to gather factual data on exposure events (e.g., whether an exposure occurred, in which department,
17 what PPE was used) rather than to measure an underlying psychological trait.

18 The second section utilized the Generalized Anxiety Disorder-7 (GAD-7) scale, a validated 7-item self-
19 report screening tool developed by Spitzer et al. .¹³ The GAD-7 is designed to assess the severity of generalized
20 anxiety disorder (GAD) symptoms and serves as an effective brief screening instrument in clinical and research
21 settings.

22 The GAD-7 scale is a 4-point Likert-type instrument (0 = not at all, 3 = nearly every day), with total
23 scores ranging from 0 to 21. The scale contains no reverse-scored items. In the evaluation of the scale, total
24 scores of 5, 10, and 15 serve as cutoff points for mild, moderate, and severe anxiety, respectively. Participants
25 with a total score of 10 or higher should undergo further diagnostic assessment to confirm the presence of
26 generalized anxiety disorder using additional methods. We categorized participants based on self-reported
27 occupational exposure to blood and body fluids, forming exposed and unexposed comparison groups for
28 subsequent analyses.

29 We assessed anxiety severity using the Turkish version of the GAD-7 scale, validated by Konkan et
30 al. .¹⁴ This adaptation demonstrated excellent internal consistency in our sample (Cronbach's $\alpha=0.89$).

1 We conducted statistical analyses using SPSS v23 for descriptive statistics (mean (SD), percentages)
2 and bivariate comparisons (Chi-square, t-test). Given significant overdispersion in GAD-7 scores
3 (variance/mean ratio = 3.7; Lagrange Multiplier test $*p* < 0.01$), multivariate analysis employed Negative
4 Binomial Regression via Python's statsmodels 0.14.0 with robust standard errors. All variables were defined as
5 follows: BBF exposure (any self-reported contact with blood or body fluids during internship, yes/no), high-risk
6 department (rotation in emergency medicine or general surgery, yes/no), age (continuous, in years), gender
7 (female/male), and contributing factors (self-reported reasons for exposure, with multiple selections allowed).
8 For the negative binomial regression model, predictors included BBF exposure status (yes/no), rotation in a
9 high-risk department (emergency medicine/general surgery; yes/no), age, gender, and self-reported
10 contributing factors to exposure (e.g., performance anxiety, pressure from assistants). Categorical predictors
11 were coded as binary variables (0 = no, 1 = yes). No categories were collapsed or treated as sparse, as each
12 contributing factor had sufficient response frequencies to be included as an individual predictor. These
13 contributing factors, while only reported by the exposed group, were included in the full model to assess their
14 association with anxiety severity across all participants, acknowledging their origin as key themes from the
15 exposed subgroup. We did not conduct a formal power analysis due to the census approach; however, the final
16 sample size of 203 participants (response rate: 74.9%) was sufficient to detect moderate effect sizes in group
17 comparisons with 80% power at $\alpha=0.05$.

18 There were no missing data in the completed questionnaires, as all items were required for electronic
19 submission. Therefore, no imputation or additional handling of missing data was necessary.

20 **Ethics approval**

21 The authors received permission from Bursa Uludağ University Health Sciences Research and
22 Publication Ethics Committee with the board decision dated 29.01.2025 and numbered 2025-01/3.

23

1 RESULTS.

2 We present the Demographic and Exposure Characteristics of the study participants in Table 1.

3 The study included 203 students, with a gender distribution of 58.6% female (n=119) and 41.4% male
4 (n=84). Age distribution analysis revealed that 35% of participants (n=71) were 24 years old, while 22.2% (n=45)
5 were 23 years old.

6 We found that 56.2% of participants (n=114) reported blood/bodily fluid exposures, with the following
7 distribution: cutaneous (23.6%, n=48), percutaneous (12.8%, n=26), and dual-route exposures (10.8%, n=22).

8 We present the Exposure context and risk factors of the study participants in Table 2 (n=114 for
9 exposure-related data).

10 16.7% of exposed interns (n=19/114) reported blood/body fluid exposures during their first internship
11 month. We observed a marked decline in exposure incidence after the first training month (from 16.7% to 10.5%,
12 $\chi^2=4.1$, $p=0.03$). We interpret this trend in the discussion section. Of the exposed participants, 32.5% (n=37/114)
13 reported a single incident, while 67.5% (n=77/114) experienced multiple exposures (≥ 2 incidents) ($\chi^2=27.67$,
14 $p<.001$). Among those with recurrent exposures, the distribution was: 29.8% of exposed participants (n=34/114)
15 had two exposures, 19.3% (n=22) three exposures, 8.8% (n=10) four exposures, and 9.6% (n=11) five or more
16 exposures.

17 Exposures were most frequently during venous blood draws (50.0%, n=57), followed by arterial
18 punctures (33.3%, n=38). When we evaluated exposure according to the department in which they performed
19 their internship, we found that 66.7% of the exposed interns experienced blood or body fluid exposures during
20 their emergency department internship, 25.4% during their general surgery internship, and 23.7% during their
21 internal medicine internship.

22 Among exposed participants, 89.5% used PPE; we found that participants universally adopted gloves
23 (100%), mask usage was 58.8%, but gown usage was low (16.7%). The most frequently cited contributing
24 factors were intense working hours (31.6%), the pressure from assistants (21.9%), and inexperience (20.2%).

25 We present the prevalence of anxiety among final-year medical students in Table 3.

26
27 Half of final-year medical students (50.2%) screened positive for anxiety. The mean GAD-7 score was
28 significantly higher among exposed students (7.21 vs. 5.39, $p=0.016$). Table 4 presents the distribution of
29 anxiety severity by exposure status. Although the distribution of anxiety severity categories did not differ
30 significantly between exposure groups ($\chi^2=3.43$, $p=0.33$), 75% of participants with severe anxiety were in the
31 exposed group. A non-significant trend toward higher anxiety prevalence was observed among exposed

1 students when combining mild-to-severe categories (54.4% vs. 45.0%; $\chi^2=2.51$, $p=0.11$; $RR=1.21$, 95% CI 0.91–
2 1.61), warranting further investigation in larger samples. We present the Negative Binomial Regression Results
3 model in Table 5.

4 Our negative binomial regression revealed that blood/body fluid exposure is associated with a 34%
5 increase in expected anxiety severity ($IRR=1.34$, 95% CI [1.18-1.52], $p<0.001$), equivalent to a 2.1-point rise in
6 mean GAD-7 scores. High-risk department rotations (emergency/surgery) amplified anxiety risk by 52%
7 ($IRR=1.52$, [1.29-1.80], $p<0.001$). Among exposure-contributing factors, performance anxiety ($IRR=1.85$, [1.32-
8 2.60]) and pressure from assistants ($IRR=1.61$, [1.27-2.04]) demonstrated stronger effects than exposure itself.
9 Notably, interns citing pre-existing anxiety as a contributor had 92% higher scores ($IRR=1.92$, [1.15-3.22]). Age
10 and gender showed no significant associations.

11

12 **DISCUSSION.**

13 Our findings suggest that more than half of the final-year medical students may have experienced
14 occupational exposure based on self report during clinical training, with varying rates that other studies have
15 reported internationally: while higher than the 20.9% incidence that Souza-Borges et al. reported in Brazil (where
16 56.2% of cases involved sharps injuries)¹⁵, it aligns closely with Inga et al.'s Peruvian cohort (51.5% reporting
17 at least one biological accident).¹⁶ However, our rates remain slightly lower than the 70.0% exposure frequency
18 that Lee's study documented.¹⁷

19 In terms of exposure routes, our study revealed that 23.6% of participants experienced cutaneous
20 contact with blood or bodily fluids, while 12.8% sustained percutaneous injuries (such as needlestick or sharps-
21 related incidents). Additionally, 10.8% of respondents reported both types of exposure. These figures are
22 consistent with Inga et al., who reported that 80.6% of biological accidents involved contact with blood, and
23 47.6% resulted from sharps injuries.¹⁶ Our findings also align with Alpat et al., who observed that 82.9% of
24 medical trainees' high-risk exposures were due to needlestick injuries, followed by mucosal (35.7%) and sharps-
25 related lacerations (30%).¹⁸ We also noted the overall frequency of exposure: while 32.5% of our study reported
26 a single incident, over two-third (67.5%) experienced repeated exposures (two or more), suggesting that
27 cumulative risk increases with ongoing clinical practice. This pattern echoes Karstaedt and Pantanowitz's
28 findings, in which 69% of interns reported at least one percutaneous exposure, and 30% recalled three or more
29 during internship.¹⁹ Such trends point to the importance of continued risk throughout training, especially in high-
30 procedure environments and underscore the need for continuous, rather than one-off, safety education within
31 medical curricula.

1 The timing of exposure appears to be closely associated with clinical inexperience. Notably, 16.7% of
2 exposed interns reported their first exposure during the initial month of internship, with subsequent rates
3 declining to 10.5% in the second and third months. This pattern suggests a possible learning curve effect, where
4 early vulnerability may stem from unfamiliarity with procedures and institutional safety protocols. Karani et al
5 similarly reported that poor clinical skills and lack of supervision were associated with the significantly higher
6 rates of accidental exposure among first-year trainees.²⁰ The fact that over two-thirds of our participants
7 experienced repeated exposures suggests a need for structured early-phase training programs and close
8 mentorship during the first months of clinical rotations. Therefore, integrating enhanced procedural simulation
9 and staged task delegation into early clinical curricula may help mitigate risk during this critical adaptation
10 period.

11 Clinical procedures associated with exposure were primarily invasive in nature. Half of all incidents
12 occurred during venous blood draws (50.0%), followed by arterial punctures (33.3%) and surgical suturing
13 (29.8%). These findings are consistent with Karani et al., who reported that 38% of interns experienced injuries
14 during phlebotomy and 19% while assisting with sutures.²⁰ The predominance of venipuncture-related injuries
15 mirrors global patterns, including WHO's identification of phlebotomy as a high-risk procedure for needlestick
16 injuries.²¹ Exposure rates also varied notably by clinical rotation, with the highest prevalence observed in
17 emergency medicine (66.7%), followed by general surgery (25.4%) and internal medicine (23.7%). This
18 distribution suggests that fast-paced, high-volume environments such as emergency departments may amplify
19 procedural risks, particularly for less-experienced trainees. Aigbodion et al. similarly observed elevated
20 exposure frequencies during high-intensity rotations, particularly in surgery and obstetrics.²²

21 Although 89.5% of exposed participants reported using PPE at the time of exposure, the type and
22 completeness of PPE varied. While nearly all students used gloves (100.0%), only 58.8% wore masks and just
23 16.7% used gowns. These rates suggest partial adherence to recommended protocols, consistent with findings
24 by Lopes et al., who noted that while most healthcare professionals wore gloves and masks, the use of
25 protective gowns, eye shields, and caps was suboptimal.²³ Inadequate PPE usage, particularly in procedures
26 involving splashing or sharps handling, may contribute to unnecessary exposure risk. Regarding contextual
27 factors, students identified extended working hours (17.7%), hierarchical pressure (12.3%), and lack of
28 experience (11.3%) as the primary contributors to exposure. These self-reported drivers align with Pereira et
29 al.'s observations of increased occupational injuries associated with both mechanical and psychosocial risk
30 factors, especially under stressful conditions.²⁴ Such findings highlight the importance of not only providing
31 adequate PPE, but also fostering a culture of safety, supervision, and psychological support within training

1 environments. While our study did not directly measure training effectiveness, these findings highlight potential
2 gaps between knowledge and practice that warrant further investigation.

3 Despite existing formal training, as evidenced by participants completing our university's mandatory
4 safety education, which includes:

- 5 • Transition to Clinic-Orientation Day (2021 curriculum): Dedicated modules on occupational
6 risks, infection control, and stress management.
- 7 • Emergency Department-specific pre-rotation orientations: Covering BBF exposure risks,
8 sharps injury prevention, and PPE protocols.

9 our data reveal persistent gaps between knowledge and practice.

10 Specifically, we observed: 56.2% exposure prevalence (higher than global averages); 66.7% exposures
11 in ER (peak risk department); 67.5% recurrent exposures (≥ 2 incidents); and critically, gown usage at only
12 16.7% during exposures.

13 Roberts (2023) emphasizes that sharps injuries persist despite safety training, particularly among early-
14 stage trainees and with non-engineered devices.²⁵ This aligns with Abernethy et al.'s (2020) report of 64% exposure
15 prevalence among trained healthcare workers.²⁶ Further supporting this theory-practice gap, Datar et al. (2022)
16 identified significant discrepancies between knowledge and practice in needlestick injury prevention among
17 medical students mirroring our findings of recurrent exposures (67.5%) and low gown usage (16.7%) despite
18 institutional training.²⁷

19 This theory-practice disconnect suggests that current training may not fully address key areas critical
20 for practical readiness and sustained safety behaviors, including:

21 Real-world stress dynamics: Performance anxiety (IRR=1.85) and hierarchical pressure (IRR=1.61)
22 persist despite didactic coverage, indicating a need for training that simulates and helps manage these
23 psychological stressors in high-stakes environments.

24 Procedural fluency and mastery: 50% of exposures occurred during venipuncture a basic skill taught
25 early suggesting that initial skill acquisition may not translate into robust, error-proof execution under pressure.

26 Behavioral sustainability and compliance: Recurrent exposures and consistently low PPE compliance
27 (e.g., gown usage) indicate a failure in the long-term retention and consistent application of safety protocols.

28 To proactively address these potential gaps and enhance practical preparedness, we propose the
29 following targeted interventions, which warrant further exploration and integration into medical curricula:

30 Immersive ER simulations: Implementing Virtual Reality (VR) scenarios that realistically replicate
31 chaotic trauma bays with bleeding patients could provide a safe environment for students to practice high-risk

1 procedures under simulated pressure, thereby improving performance anxiety and procedural fluency. These
2 interventions require further evaluation in future studies.

3 "Stress-tested" PPE drills: Conducting unannounced mock exposures or rapid-response PPE drills
4 during clinical shifts could enhance muscle memory and ensure consistent, correct PPE usage under realistic
5 conditions, addressing behavioral sustainability.

6 Anxiety-inoculation through integrated CBT techniques: Embedding cognitive-behavioral therapy (CBT)
7 techniques from existing stress management modules directly into high-risk procedure training could help
8 students recognize and manage their anxiety in real-time, preventing the stress-injury cycle.

9 Table 3 shows that 50.2% of final-year medical students in our study screened positive for anxiety, with
10 7.9% meeting the criteria for severe symptoms. This prevalence substantially exceeds the global average of
11 33.8% for medical students, as Quek et al. reported in a recent meta-analysis, and is also markedly higher than
12 general population estimates, which range from 3% to 25%.^{28,29} Moreover, we observed a statistically significant
13 difference in GAD-7 scores between participants with and without exposure to blood or body fluids: the exposed
14 group reported a mean score of 7.21 (5.38), significantly higher than the unexposed group [5.39 (5.15); $p=0.016$]
15 (Table 4). Although the difference in anxiety severity categories (normal, mild, moderate, severe) did not reach
16 statistical significance ($p=0.33$), 75% of those with severe anxiety had experienced exposure (Table 5),
17 suggesting a vulnerable subgroup.

18 Although the non-significant chi-square test for categorical anxiety severity and the significant findings
19 from the t-test and regression model may seem contradictory, this reflects the different nature of categorical
20 and continuous variables. Categorizing continuous data into ordinal groups results in loss of information and
21 reduced statistical efficiency³⁰. The continuous GAD-7 scores therefore provide a more sensitive assessment
22 of the exposure-anxiety relationship

23 This finding aligns with prior research: Gaspar et al. demonstrated that workplace injuries can
24 exacerbate anxiety, forming a "stress-injury" feedback loop.¹¹ Similarly, Granger and Turner's meta-analysis
25 supports a bidirectional relationship between occupational trauma and anxiety disorders in healthcare
26 workers.¹² These findings highlight the potential need for integrated occupational and psychological safety
27 protocols in medical education, though further multi-center studies are needed to confirm generalizability.

28 In our study psychological factors, particularly hierarchical pressure (IRR=1.61) and performance
29 anxiety (IRR=1.85), exerted stronger effects on anxiety than physical exposure (IRR=1.34). This suggests that
30 the culture of medical training may pose greater mental health risks than occupational hazards themselves. The
31 92% increase in anxiety among interns who identified 'anxiety level' as an exposure contributor (IRR=1.92) is

1 consistent with the conceptual framework of the stress-injury cycle proposed by Gaspar et al. (2020), though
2 our cross-sectional design cannot establish causality.¹¹

3 The findings of this study underscore the urgent need to enhance occupational safety education and
4 psychological support within undergraduate medical curricula. The high rate of exposure, particularly during
5 early internship months and in high-intensity departments like emergency medicine, signals a critical gap in
6 procedural preparedness and supervision. Incorporating structured simulation-based training, task-specific risk
7 briefings, and progressive skill acquisition modules may help reduce preventable exposures. Additionally,
8 institutional policies should foster a non-punitive culture of reporting and ensure full compliance with PPE
9 protocols. From a mental health perspective, the association between exposure and elevated anxiety highlights
10 the necessity of integrating routine psychological screening and peer support systems into internship programs.
11 Aligning with the principles outlined in the EU Directive 2010/32, interventions should be proactive,
12 comprehensive, and culturally tailored to mitigate both physical and emotional harm.³⁰ Ultimately, fostering a
13 culture of safety from the earliest stages of clinical training may yield long-term benefits for both patient care
14 and workforce well-being.

15 **LIMITATIONS**

16 This study has several limitations that we acknowledge. First, the cross-sectional design fundamentally
17 limits the strength of causal inferences between blood/body fluid exposures and anxiety levels, preventing
18 conclusions about direct causation or directionality of effect. This inherent limitation may also contribute to the
19 observed modest effect sizes and the lack of statistical significance in certain key associations, such as anxiety
20 severity categories, as highlighted by our findings.

21 Second, reliance on self-reported data introduces potential recall bias (particularly for exposure
22 incidents occurring early in internship) and social desirability bias in PPE compliance reporting. This self-
23 reported nature, coupled with the absence of objective exposure verification and a lack of triangulation methods
24 (e.g., direct observation, institutional records), limits data robustness and the ability to fully ascertain the true
25 incidence and impact of exposures.

26 Third, the single-center nature of this study at a medical school constrains generalizability to other
27 cultural or healthcare contexts.

28 Fourth, the 25.1% non-response rate may affect representativeness, as students with high anxiety might
29 have been less likely to participate.

1 Fifth, while the GAD-7, although validated for screening, it cannot establish clinical diagnoses of anxiety
2 disorders. Finally, related to the second point, the absence of objective exposure verification could lead to
3 underreporting or inaccurate reporting of exposure frequency and severity.

4 Sixth, while we used the GAD-7 as a validated screening tool for anxiety symptoms, it does not provide
5 a clinical diagnosis of anxiety disorders. Additionally, we did not assess other psychological factors such as
6 depression, burnout, coping styles, or prior psychiatric history, nor did we measure potential confounders such
7 as workload intensity, sleep patterns, caffeine use, or prior occupational safety training, all of which may
8 influence both anxiety levels and exposure risk. Therefore, the stress-injury cycle framework, while conceptually
9 relevant, remains hypothetical in the context of our study and requires further investigation with longitudinal
10 designs. Despite these limitations, our findings align with global evidence on occupational risks in medical
11 training and provide actionable insights for educational reform.

12 Therefore, our prevalence estimates should be interpreted with caution, and the findings regarding
13 exposure frequency and timing may be subject to recall error.

14 **CONCLUSION**

15 This study found that over half of final-year medical students (56.2%) reported exposure to blood or
16 body fluids during clinical training, with multiple incidents reported by 67.5%. Venous blood draws, emergency
17 department rotations, and early internship months emerged as key risk contexts. Despite comprehensive safety
18 training (including 2021 curriculum modules and ER-specific orientations), incomplete PPE adherence (gown
19 use: 16.7%) and contributing factors like hierarchical pressure (21.9%) persisted. We found that exposure was
20 significantly associated with elevated anxiety scores (mean GAD-7: 7.21 vs. 5.39, $p=0.016$), indicating a dual
21 physical-mental health burden. These findings suggest a need to further examine and potentially adapt existing
22 training approaches to better integrate stress-adapted simulations, real-time compliance feedback, and
23 psychological safety practices. These findings generate hypotheses about potential curricular improvements
24 and suggest the need for further research with objective measures of training effectiveness.

25 **STATEMENTS AND DECLARATIONS**

26 **Acknowledgments:** None

27 **Funding/Support:** This research received no specific grant from any funding agency.

28 **Other disclosures:** None

1 **Ethical approval:** The authors received permission from Bursa Uludağ University Health Sciences
2 Research and Publication Ethics Committee with the board decision dated 29.01.2025 and numbered 2025-
3 01/3.

4 **Disclaimers:** None

5 **Previous presentations:** None

6 **Data:** In this study, no data from outside sources were used. All data were collected directly from
7 participants after ethical approval was obtained.

8 **Consent to participate:** Written informed consent was obtained from all participants prior to enrollment.
9 The online questionnaire commenced with a detailed Informed Voluntary Consent Form outlining the study
10 objectives, procedures, potential risks/benefits, and data confidentiality measures. Participants were informed
11 that their participation was voluntary and that they could withdraw at any time without penalty. Only those who
12 electronically acknowledged their consent proceeded to the study questions.

13 **Consent for publication:** Not applicable. The manuscript contains no individual person's data.

14 Declaration of conflicting interest: The authors declare that there is no conflict of interest.

15 **Data availability:** The anonymized dataset is available from the corresponding author upon reasonable
16 request.

17

18

19

Accepted, in press

1 SUMMARY - ACCELERATING TRANSLATION

2 **Title:** Medical Students' Risk of Exposure to Blood and Body Fluids and Its Impact on Anxiety Levels

3 **The Main Problem to Solve**

4 Final-year medical students (interns) who are in clinical training in hospitals are at risk of coming into contact
5 with patients' blood or other body fluids (e.g., saliva, urine). This can happen through needlestick injuries, cuts,
6 or splashes. Such exposures pose a **serious health risk**, including the potential transmission of blood-borne
7 diseases like Hepatitis or HIV.

8 However, the risk is not just physical. Experiencing such an incident can cause **intense stress and anxiety** for
9 students. The worry of "Did I contract a disease?" can negatively affect a student's psychological well-being
10 and, consequently, their education and professional performance. It is a critical question whether current safety
11 training programs are effectively preventing this dual (physical and psychological) risk.

12 **Aim of the Study**

13 The primary goal of this study was to determine how frequently final-year medical students are exposed to blood
14 and body fluids during their internships. Secondly, we aimed to scientifically investigate whether experiencing
15 such an exposure increases students' anxiety levels.

16 **Methodology**

17 We conducted this research with the participation of 203 medical interns. An online questionnaire was sent to
18 the participants, which consisted of three parts:

- 19 1. **Demographic Information:** Basic information such as age and gender.
- 20 2. **Exposure History:** We asked questions like, "Have you ever had a needlestick injury or been splashed
21 with blood during your internships?", "In which department did this occur?", and "What protective
22 equipment were you using at the time?"
- 23 3. **Anxiety Level Scale:** We used the **GAD-7 (Generalized Anxiety Disorder-7)** scale, a widely used
24 and reliable test, to measure the participants' anxiety levels over the past two weeks.

25 We analyzed the collected data using statistical methods to examine the relationship between exposure to blood
26 and body fluids and anxiety levels.

27 **Results**

28 Our study yielded several noteworthy findings:

- 29 • **High Rate of Exposure:** More than one in two students (56.2%) reported being exposed to blood or
30 body fluids at least once during their internships. Of those exposed, 67.5% experienced multiple
31 incidents.
 - 32 • **The Emergency Department is the Riskiest:** The most common setting for these incidents was the
33 Emergency Department (66.7%).
 - 34 • **Deficiencies in Protective Equipment Use:** While all students reported using gloves during the
35 incident, the use of gowns was critically low (16.7%).
 - 36 • **The Link Between Exposure and Anxiety:** Our most significant finding was that the anxiety scores of
37 exposed students were statistically higher than those of non-exposed students (7.21 vs. 5.39).
 - 38 • **Other Factors Increasing Anxiety:** Our analysis also showed that, in addition to exposure, working in
39 high-risk departments and the fear of making mistakes during procedures (performance anxiety)
40 significantly increased anxiety levels.
- 41

1 **REFERENCES.**

- 2 1. Azap A, Ergönül O, Memikoğlu KO, Yeşilkaya A, Altunsoy A, Bozkurt GY, et al. Occupational exposure to
3 blood and body fluids among health care workers in Ankara, Turkey. *Am J Infect Control*. 2005;33(1):48-
4 52.
- 5 2. Puro V, De Carli G, Cicalini S, Soldani F, Balslev U; European Occupational Post-Exposure Prophylaxis
6 Study Group. European recommendations for the management of healthcare workers occupationally
7 exposed to hepatitis B virus and hepatitis C virus. *Euro Surveill*. 2005;10:260-264.
- 8 3. Prüss-Ustün A, Rapiti E, Hutin Y. Estimation of the global burden of disease attributable to contaminated
9 sharps injuries among health-care workers. *Am J Ind Med*. 2005;48(6):482-490.
- 10 4. Osborn EH, Papadakis MA, Gerberding JL. Occupational exposures to body fluids among medical students:
11 a seven-year longitudinal study. *Ann Intern Med*. 1999;130(1):45-51.
- 12 5. Triassi M, Pennino F. Infectious risk for healthcare workers: evaluation and prevention. *Ann Ig*. 2018;30(4
13 Suppl 1):48-51.
- 14 6. Zhang L, Ai Y, Liu J, Yue N, Xuan J, Bal V, et al. Economic burden of needle-stick injury among healthcare
15 workers in China. *Value Health*. 2018;21:S86.
- 16 7. Erdem Y, Talas MS. Blunt and penetrating object injuries in housekeepers working in a Turkish university
17 hospital. *Am J Infect Control*. 2006;34(4):208-214.
- 18 8. Gücük M, Karabey S, Yolsal N, Özden YI. İstanbul Tıp Fakültesi Genel Cerrahi Kliniği çalışanlarında kesici-
19 delici alet yaralanmaları. *Hastane Enfeksiyonları Dergisi*. 2002;6(2):72-81.
- 20 9. Kışioğlu AN, Öztürk M, Uskun E, Kırbıyık S. Bir üniversite hastanesi sağlık personelinde kesici-delici
21 yaralanma epidemiyolojisi ve korunmaya yönelik tutum ve davranışlar. *Türkiye Klinikleri J Med Sci*.
22 2002;22:390-396.
- 23 10. Moon CS, Hwang JH, Lee CS, Park KH, Kim ES. Exposure to blood and body fluid among medical students
24 in Korea. *Am J Infect Control*. 2010;38(7):582-583.
- 25 11. Gaspar FW, Jolivet DN, Wizner K, Schott F, Dewa CS. Pre-existing and new-onset depression and anxiety
26 among workers with injury or illness work leaves. *J Occup Environ Med*. 2020;62(10):e567-e572.
- 27 12. Granger S, Turner N. Work injuries and mental health challenges: a meta-analysis of the bidirectional
28 relationship. *Pers Psychol*. 2025;78(1):11-57.
- 29 13. Spitzer RL, Kroenke K, Williams JBW, Löwe B. A brief measure for assessing generalized anxiety disorder:
30 the GAD-7. *Arch Intern Med*. 2006;166(10):1092-1097.
- 31 14. Konkan R, Şenormancı Ö, Güçlü O, Aydın E, Sungur MZ. Yaygın anksiyete bozukluğu-7 (YAB-7) testi
32 Türkçe uyarlaması, geçerlik ve güvenilirliği. *Noropsikiyatri Ars*. 2013;50(1):53-58.
- 33 15. Souza-Borges FRF, Ribeiro LA, Oliveira LCM. Occupational exposures to body fluids and behaviors
34 regarding their prevention and post-exposure among medical and nursing students at a Brazilian public
35 university. *Rev Inst Med Trop Sao Paulo*. 2014;56(2):157-163.
- 36 16. Inga E, López G, Kamiya C. Accidentes biológicos en estudiantes de medicina de una universidad peruana:
37 prevalencia, mecanismos y factores de riesgo. *An Fac Med (Lima)*. 2011;71(1):37.
- 38 17. Lee CS, Hwang JH, Seon SY, Jung MH. Exposure to blood and body fluids for medical students during
39 clerkship. *Korean J Med*. 2008;74(5):500-505.
- 40 18. Alpat S, Özgüneş I, Nemli S, Erben N. Tıp fakültesi öğrencilerinde kan ve vücut sıvılarıyla temasa bağlı
41 mesleki enfeksiyon riski. *Flora*. 2008;13(2):88-92.

- 1 19. Karstaedt AS, Pantanowitz L. Occupational exposure of interns to blood in an area of high HIV
2 seroprevalence. *S Afr Med J.* 2001;91(1):57-61.
- 3 20. Karani H, Rangiah S, Ross AJ. Occupational exposure to blood-borne or body fluid pathogens among
4 medical interns at Addington Hospital, Durban. *S Afr Fam Pract.* 2011;53(5):462-466.
- 5 21. Cadman H, editor. *WHO guidelines on drawing blood: best practices in phlebotomy.* Geneva: WHO Press;
6 2010.
- 7 22. Aigbodion SJ, Motara F, Laher AE. Occupational blood and body fluid exposures and human
8 immunodeficiency virus post-exposure prophylaxis amongst intern doctors. *S Afr J HIV Med.*
9 2019;20(1):958.
- 10 23. Lopes TL, Pinheiro AF, Silva CG, Mocelin PC, Zanin GD. Utilização, opinião e conhecimento dos
11 profissionais de saúde sobre equipamentos de proteção individual durante a pandemia de SARS-CoV-2:
12 um estudo transversal. *Res Soc Dev.* 2021;10(12):e26101220016.
- 13 24. Pereira A, Ribeiro A, Miranda G, Silva P, Silva A, Soares J, et al. Caracterização dos acidentes de trabalho
14 num hospital terciário português entre 2017 e 2022. *Rev Port Saude Ocup.* 2023;15.
- 15 25. Roberts LW. Working together to ensure greater safety in our work and training environments. *Acad Med.*
16 2023;98(7):755-756.
- 17 26. Abere G, Yenealem DG, Wami SD. Occupational exposure to blood and body fluids among health care
18 workers in Gondar Town, Northwest Ethiopia: a cross-sectional study. *J Environ Public Health.*
19 2020;2020:3640247.
- 20 27. Datar UV, Kamat M, Khairnar M, Wadgave U, Desai KM. Needlestick and sharps injury in healthcare
21 students: prevalence, knowledge, attitude and practice. *J Family Med Prim Care.* 2022;11(10):6327-6333.
- 22 28. Quek TTC, Tam WWS, Tran BX, Zhang M, Zhang Z, Ho CSH, et al. The global prevalence of anxiety among
23 medical students: a meta-analysis. *Int J Environ Res Public Health.* 2019;16(15).
- 24 29. Baxter AJ, Scott KM, Vos T, Whiteford HA. Global prevalence of anxiety disorders: a systematic review and
25 meta-regression. *Psychol Med.* 2013;43(5):897-910.
- 26 30. Zhao LP, Kolonel LN. Efficiency loss from categorizing quantitative exposures into qualitative exposures in
27 case-control studies. *Am J Epidemiol.* 1992;136:464-474.
- 28 31. Yaneva-Deliverska M. European Union regulation on prevention of sharps injuries in hospital and healthcare
29 sector. *J IMAB.* 2012;18(3):357-359.
- 30

1 **TABLES**

2

3 **Table 1.** Demographic and Exposure Characteristics (N=203)

4

Characteristic	Category	n	%
Gender	Women	119	58.6
	Men	84	41.4
Age (years)	22	7	3.4
	23	45	22.2
	24	71	35.0
	25	39	19.2
	26	18	8.9
	≥27	23	11.3
Any exposure	Yes	114	56.2
	No	89	43.8
Exposure route* (n=114)	Cutaneous only	48	23.6
	Percutaneous injury (CPI)	26	12.8
	Mucosal only	3	1.5
	CPI + Cutaneous	22	10.8
	CPI + Mucosal + Cutaneous	7	3.4
	CPI + Mucosal	6	3.0
	Mucosal + Cutaneous	2	1.0

5 *Multiple selections were made

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

1 **Table 2.** Exposure Context and Risk Factors (n=114 for exposure-related data)*
2

Characteristic	Category	n	%
Training month at first exposure	Month 1	19	16.7
	Month 2	12	10.5
	Month 3	12	10.5
Exposure frequency	1 time	37	32.5
	2 times	34	29.8
	3 times	22	19.3
	4 times	10	8.8
	≥5 times	11	9.6
Procedures associated with exposure*	Venous blood draw	57	50.0
	Arterial puncture	38	33.3
	Surgical suturing	34	29.8
Clinical rotations associated with exposure*	Emergency Department	76	66.7
	General Surgery	29	25.4
	Internal Medicine	27	23.7
PPE (Personal Protective Equipment) use during exposure	Yes	102	89.5
	No	11	9.6
PPE components*	Gloves	102	100
	Mask	60	58.8
	Gown	17	16.7
Reported Most Contributing Factors	Intense working hours	36	31.6
	Pressure from assistants	25	21.9
	Inexperience	23	20.2
	Occupational fatigue	17	14.9
	Time pressure	7	6.1
	Performance anxiety	3	2.6
	High anxiety	2	1.8

3 *Multiple selections were allowed for procedures, clinical rotations, PPE components, and contributing factors.
4 All percentages in Table 2 are based on the exposed subgroup (n=114) and represent the proportion of exposed
5 participants who reported each item. Percentages may sum to more than 100% due to multiple selections.
6

1 **Table 3.** The Prevalence of Anxiety Among Final-Year Medical Students (N=203)

Severity Level	n	%	95% CI
Mild	60	29.2	23.1-35.9
Moderate	26	12.8	8.6-18.2
Severe	16	7.9	4.6-12.4
Total with anxiety	102	50.2	43.3-57.1

2
3

Accepted, in-press

1 **Table 4.** Anxiety Severity by Blood/Body Fluid Exposure Status (N=203)

Anxiety Severity	Exposed		Unexposed		Total
	n	%	n	%	n
Normal	52	51.5	49	48.5	101
Mild	35	58.3	25	41.7	60
Moderate	15	57.7	11	42.3	26
Severe	12	75.0	4	25.0	16
Total	114	56.2	89	43.8	203

2
3
4

We present the Negative Binomial Regression Results model in Table 5.

Accepted, in-press

1 **Table 5.** Multivariate Negative Binomial Regression for Anxiety Severity (GAD-7 Scores)

Predictor	IRR	95% CI	p-value
Blood/Body Fluid Exposure	1.34	[1.18, 1.52]	<0.01**
High-Risk Department*	1.52	[1.29, 1.80]	<0.01**
Contributing Factors (Ref: Inexperience)			
Fatigue	1.18	[0.94, 1.48]	0.15
Long Working Hours	1.42	[1.13, 1.78]	0.01**
Pressure from Assistants	1.61	[1.27, 2.04]	<0.01**
Performance Anxiety	1.85	[1.32, 2.60]	<0.01**
Time Management Issues	0.97	[0.72, 1.31]	0.84
Anxiety Level	1.92	[1.15, 3.22]	0.01*
Age	0.99	[0.97, 1.02]	0.54
Gender (Female)	0.98	[0.83, 1.16]	0.82

2 *High-risk departments: Emergency Medicine and General Surgery.

3 † These contributing factors were reported by participants in the exposed group (n=114) and were included in
 4 the full model (N=203) to assess their association with anxiety severity. Model Fit: Log-likelihood = -438.7, AIC
 5 = 901.4; **p<0.01, *p<0.05

Accepted, in